



CITY OF KWINANA
WESTERN AUSTRALIA

BLACK COCKATOO CONSERVATION ACTION PLAN



Acknowledgment of Country

We acknowledge the Traditional Custodians of the boodjar (lands) on which this plan was prepared, the Whadjuk and Bindjareb Noongar¹ people, including Elders past and present.

We respect the enduring connections of Noongar people to the southwest of Western Australia, which includes the last remaining habitat for southwest Western Australia's threatened black cockatoos.

Please note: this document may contain the words and names of people who have passed away.



¹ There are several ways to spell Noongar; as is common for oral, non-written languages. This plan uses 'Noongar', and acknowledges the diversity of spellings.



BLACK COCKATOO CONSERVATION ACTION PLAN

Keep Carnaby's Flying – Ngoolarks Forever



BLACK COCKATOO CONSERVATION

Suggested citation

Murdoch University Black Cockatoo Conservation Management Project (2024) *City of Kwinana Black Cockatoo Conservation Action Plan, Keep Carnaby's Flying – Ngoolarks Forever* project, Murdoch University.

COVER PHOTO Rick Dawson

PHOTO CREDITS Karen Riley, Sam Rycken, Molly Spaulding, Zoe Kissane (page 18 and page 44) (Murdoch University black cockatoo researchers)

Partners

This plan was developed for the City of Kwinana by Murdoch University as part of the City's participation in the *Keep Carnaby's Flying – Ngoolarks² Forever* project, a community activation initiative to help safeguard black cockatoos in the Perth-Peel region, made possible with support from Lotterywest and the following partner organisations:



2 There are several ways to spell and say the Noongar name for Carnaby's cockatoo. This plan uses 'ngoolark', and acknowledges the name diversity.

Contents

Acknowledgment of Country	II	Threat 2. Lack of fresh drinking water	37
Suggested citation	IV	2.1 Heat stress.....	37
Partners	IV	2.2 Poor water quality.....	38
Introduction	4	2.3 Vehicle strike when drinking on roads and verges.....	38
Purpose	4	Threat 3. Vehicle strike	41
Background	4	Threat 4. Carnaby’s Hindlimb Paralysis Syndrome	44
Murdoch University’s black cockatoo research team – Who we are	5	Threat 5. Raven attack	45
Southwest Western Australia’s black cockatoos at a glance	6	Threat 6. Illegal shooting	46
Cultural significance of black cockatoos	8	Threat 7. Climate change and Urban Heat Island effects	47
Importance of the City of Kwinana for black cockatoos	10	Threat 8. Cumulative impacts	50
Threat 1. Loss of habitat	12	Actions table: Summary of actions to safeguard black cockatoos in the City of Kwinana	51
1.1 Lack of food resources.....	12	Appendices	54
1.2 Lack of roosts.....	29	Appendix 1. Selected references and resources..	54
1.3 Lack of vegetation corridors.....	32	Appendix 2. Black cockatoos from the City of Kwinana: Causes of admission to Perth Zoo Veterinary Department 2018–23.....	56
1.4 Lack of breeding habitat.....	34		
1.5 Tree diseases.....	36		
1.6 Habitat loss from fire.....	36		



BLACK COCKATOO CONSERVATION



A note on referencing:

The information in this Plan about specific threats to black cockatoos and effective conservation actions comes from peer-reviewed scientific publications, and from guidelines and advice prepared by government and organisations with specific black cockatoo expertise. The Plan incorporates information from ongoing research by Murdoch University and BirdLife Australia, and includes data specific to the City of Kwinana. For readability, referencing has been confined to footnotes supporting selected statements. A list of selected references and resources is provided at the end of the document. For general advice about black cockatoos, please contact BirdLife Australia.


List of figures

Figure 1. Distribution of a: Carnaby's cockatoos; b: Baudin's cockatoos; c: forest red-tailed black cockatoo.....	7
Figure 2. Known black cockatoo roost sites within the City of Kwinana	10
Figure 3. GPS data showing foraging movements for a tagged Carnaby's cockatoo and its flock.....	13
Figure 4. GPS data showing foraging movements for a tagged Carnaby's cockatoo and its flock.....	13
Figure 5. City of Kwinana showing council-managed green spaces in relation to known night roosts for black cockatoos	21
Figure 6. City of Kwinana: Bush Forever areas in relation to known roost sites for black cockatoo flocks, showing areas of value for the City to investigate for revegetation activities and infill planting with black cockatoo food species	22
Figure 7. City of Kwinana: DBCA reserves and legislated lands in relation to known roost sites for black cockatoo flocks.....	23

Figure 8. Schools and their ovals often provide valuable foraging and roosting habitat for black cockatoos	25
Figure 9. Recommended priority areas for habitat retention and revegetation in the City of Kwinana.....	26
Figure 10. Locations of confirmed Carnaby's cockatoo breeding activity (12km buffers)...	34
Figure 11. Recommended locations to install Cockitrough water stations.....	40
Figure 12. City of Kwinana black cockatoo admissions to Perth Zoo Veterinary Department 2018–23: Reported vehicle strikes.....	41
Figure 13. Reported locations of vehicle strike cases in the City of Kwinana, for black cockatoos admitted to Perth Zoo Veterinary Department 2018–23.....	43
Figure 14. Reported locations of shooting cases in the City of Kwinana, for black cockatoos admitted to Perth Zoo Veterinary Department 2018–23	46

List of tables

Table 1. High priority food plants to plant for black cockatoos in the City of Kwinana	19
--	----



Introduction

Black cockatoos are iconic wildlife in the Western Australian landscape. They are charismatic species that generate high levels of public attention and concern, and consistently poll amongst Australia’s favourite birds.

All three of southwest Western Australia’s endemic black cockatoo species – Carnaby’s cockatoo, Baudin’s cockatoo and forest red-tailed black cockatoo – are experiencing severe population declines due to habitat loss and other threatening processes, and face extinction within decades unless threats are addressed³. In the Perth-Peel region in particular, they are under great pressure.

As land managers and decision-makers, local governments in Perth and Peel play a critical role in protecting black cockatoos in their suburbs. Benefits of retaining black cockatoos in urban environments are numerous and include social, cultural, environmental and biodiversity benefits. In the Perth-Peel region, poor planning and management have already led to the loss of black cockatoos entirely from some areas.

The City of Kwinana is an important area for black cockatoos in the Perth-Peel region, with roost sites and food resources that support large flocks of Carnaby’s cockatoos and forest red-tailed black cockatoos⁴. For the City to retain black

cockatoos in its urban landscape, there are some specific, evidence-informed actions that it will be important for the City to undertake when managing its built and natural environments. This plan presents those actions.

Purpose

This plan outlines the actions that the City of Kwinana can undertake, to manage the key threatening processes affecting black cockatoos within the City of Kwinana. The plan has been designed as a practical guide to help the City of Kwinana safeguard its black cockatoos.

Background

In 2023, in recognition of the City of Kwinana’s importance for black cockatoos, the City was one of four local government authorities (LGAs) invited to participate in the second year of the *Keep Carnaby’s Flying – Ngoolarks Forever* project⁵, with project activities commencing in 2024. The City accepted the invitation⁶.

The *Keep Carnaby’s Flying – Ngoolarks Forever* project is a collaboration between universities, Aboriginal organisations and natural resource management and wildlife NGOs, with support from Lotterywest. The project’s goal is to activate

the community to ‘keep Carnaby’s flying’ in the Perth-Peel region, through a range of on-ground activities underpinned by science and cultural engagement. Key activities include outreach to community groups and schools, and funds and support to participating LGAs to plant black cockatoo food trees and install bird water drinking stations in targeted locations, informed by research, to help black cockatoos in the short term. Each participating LGA also receives an LGA-specific Black Cockatoo Conservation Action Plan (this document) to guide longer-term actions.

This Conservation Action Plan was developed by one of the lead organisations in *Keep Carnaby’s Flying – Ngoolarks Forever*; Murdoch University. The university has been researching southwest WA’s three black cockatoo species since 2007, with a focus on health, habitat use and flock movements, including GPS and ARGOS satellite tracking of flocks in the Perth-Peel region and across the species’ ranges (see blackcockatooconservationwa.com). This research, along with locality-specific black cockatoo data from BirdLife Australia, has enabled the development of a targeted plan for the City of Kwinana.

The plan covers:

- 1 The importance of the City of Kwinana for black cockatoos.**
- 2 Threats faced by black cockatoos in the City of Kwinana.**
- 3 Actions for the City to take to protect its black cockatoos.⁷**

³ As the International Union for Conservation of Nature (IUCN) notes in its ‘Justification of Red List category’ for Baudin’s cockatoo: “This species is experiencing very rapid declines exceeding a rate of 90% over the past three generations...There is no reason to suggest these declines will cease with nest hollow shortages continuing to afflict the species with low productivity, and fire and drought impacts [on habitat] projected to worsen with ongoing anthropogenic climate change. With a relatively small population (likely comprising fewer than 4,000 mature individuals), future declines of this magnitude place Baudin’s Black-cockatoo at considerable extinction risk; it is therefore listed as Critically Endangered.” Critically Endangered is the last step before extinction.

⁴ The southwest part of the City of Kwinana is also within the ‘likely’ distribution for Baudin’s cockatoos.

⁵ keepcarnabysflying.org.au

⁶ City of Kwinana Letter of Participation, Manager Environmental and Health Services, 7 August 2023.

⁷ While the overall *Keep Carnaby’s Flying – Ngoolarks Forever* project focused on Carnaby’s cockatoos due to the severe pressure on this species in urban areas, this plan has been designed to support all black cockatoos in the City of Kwinana, irrespective of species.

Murdoch University's black cockatoo research team – Who we are

This plan was prepared by Murdoch University, a partner organisation in *Keep Carnaby's Flying – Ngoolarks Forever*, with contributions from the university's black cockatoo conservation management research team⁸. Our team is undertaking one of the largest and longest studies of wild cockatoos in the world.

Our research focuses on the health, demographics and ecology of Western Australia's threatened black cockatoos. We have collaborations with the Department of Biodiversity, Conservation and Attractions (DBCA), Perth Zoo, Kaarakin Black Cockatoo Conservation Centre, WA Museum, World Wildlife Fund, BirdLife Australia and industry partners. Our research addresses most of the key Actions and Objectives in the national [Recovery Plans](#) for all three of southwest Western Australian's black cockatoo species.

Since 2007 we have undertaken disease screening of injured black cockatoos treated at Perth Zoo for rehabilitation back to the wild, and health monitoring of Carnaby's cockatoo nestlings throughout the species' breeding range.

In an exciting first for Australian cockatoos, we have developed the methodology to attach GPS and ARGOS satellite tags to rehabilitated black cockatoos prior to release back into wild flocks. This allows us to track flocks and identify key areas and habitat for foraging, roosting and breeding, as well as migratory movements, about which much remains unknown. We also attach telemetry tags to black cockatoos at key breeding sites to investigate habitat use at these sites.

Our tracking of flocks in the Perth-Peel region helps to identify important habitat and address threatening processes in different localities. Working with local governments, natural resource management and urban land-care NGOs and BirdLife Australia, we can identify important foraging habitat and roost trees to retain, priority locations for planting black cockatoo food species and providing bird water drinking stations, and hotspots for threats such as vehicle strikes, so that these can be managed.

High resolution version of this plan

A high resolution electronic version of this Conservation Action Plan has been provided to the City of Kwinana by Murdoch University's black cockatoo research team. The high resolution version will allow users to 'zoom in' to particular areas on the maps, to support planning for conservation actions.



Southwest Western Australia's black cockatoos at a glance



Carnaby's cockatoos



Baudin's cockatoos



Forest red-tailed black cockatoos

Distinguishing features

- » **Carnaby's cockatoos (*Zanda latirostris*), Baudin's cockatoos (*Zanda baudinii*):** white panels on tail, Carnaby's cockatoos have a shorter upper bill (beak) than Baudin's cockatoos, and are more prevalent in urban areas
- » **Forest red-tailed black cockatoos (*Calyptorhynchus banksii naso*⁹):** red panels on tail for male, red, orange and yellow on tail for females and juveniles

Public image

- » Highly visible, charismatic wildlife, including in urban landscapes
- » Intelligent, curious, social
- » Strong cultural connections for Noongar people and wider community
- » Attract community and media attention and concern

Life history

- » Long-lived, slow-breeding
- » Flocks of generally <100 birds; may be larger, especially for Carnaby's cockatoos
- » Pairs likely mate for life; breed once per year (if breeding resources are available); usually raise one fledgling

Threat status

- » **Current population trends:** decreasing
- » All three species receive special protection as Matters of National Environmental Significance under the *Environmental Protection and Biodiversity Conservation Act 1999* (EPBC Act)
- » All three species remain in grave danger of ongoing and catastrophic population decline
- » **Carnaby's cockatoo:** Listed as Endangered under *Biodiversity Conservation Act 2016* (BC Act), EPBC Act (1999) and IUCN Red List for Threatened Species (IUCN Red List); (rapid population decline of >50%–79% in 45 years¹⁰)
- » **Baudin's cockatoo:** Endangered under BC Act and EPBC Act; Critically Endangered IUCN Red List (very rapid population decline of >90% in 45 years¹¹)
- » **Forest red-tailed black cockatoo:** Vulnerable⁹ under BC Act and EPBC Act
- » **Major threats:** ongoing habitat loss and degradation; urban and industrial expansion; fires; disease; vehicle strike; displacement from hollows by competing species. Most threats now exacerbated by harmful climate changes.

Diet

- » Preferences vary by species; predominantly seeds (also flowers, nectar, grubs) from native trees, especially banksia, hakea, marri and jarrah; some introduced plants e.g. plantation pines (Carnaby's cockatoos). Food resources must be supported by roosting habitat and water in close proximity.

⁹ Taxonomic research published in 2023 argued that forest red-tailed black cockatoos, which had been considered a sub-species, should be considered a new species (Saunders & Pickup 2023). This proposed new status, if broadly accepted, may support raising the birds' conservation status to 'Endangered', given there are fewer forest red-tailed black cockatoos left than Carnaby's cockatoos, and the latter is already recognised as endangered.

¹⁰ BirdLife International (2022). *Zanda latirostris*. The IUCN Red List of Threatened Species 2022: e.T22684733A212974328. <https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T22684733A212974328.en>

¹¹ BirdLife International (2023). Species factsheet: *Zanda baudinii*. <http://datazone.birdlife.org/species/factsheet/audins-black-cockatoo-zanda-baudinii>






Figure 1. Distribution of a: Carnaby's cockatoos; b: Baudin's cockatoos; c: forest red-tailed black cockatoo

Source: Department of Sustainability, Environment, Water, Population and Communities

Distribution, movements and habitat

- » **Distribution:** Occur only in southwest WA (see **Figure 1**)
- » **Breeding seasons vary by species:** Carnaby's cockatoo and Baudin's cockatoo generally undertake breeding activities from August to February; FRTB cockatoos around October-November, and sometimes in autumn if rains and food are sufficient.
- » **Breeding movements:** Most Carnaby's cockatoo flocks spend the non-breeding season foraging in coastal areas, including the Swan Coastal Plain, to build body condition for breeding, and then migrate inland to semi-arid Wheatbelt breeding areas. Some breeding occurs closer to the coast. Baudin's cockatoos and FRTB cockatoos breed in forests in humid and sub-humid zones; some flocks undertake seasonal movements for breeding, while others breed in their local area.
- » **Nesting:** All species nest in large hollows in large old eucalypts, living and dead, in forests and woodlands as well as remnant patches. Trees with suitable hollows are typically >200 years old. Breeding habitat requires sufficient foraging habitat and water within close proximity.
- » **Night roost habitat:** preferred roost trees are tall (often >25m), native or non-native, with medium foliage density¹². Several stands of roost trees within a few hundred metres may form a single roost site; different flocks may share a roost. Roosts must be supported by proximity to water and food; number of birds at a roost correlates with nearby food availability.
- » **Daily foraging movements:** While resident¹³ at a roost, flocks generally forage within 4–6km from that roost.



The information presented here about roosting habitat and daily foraging distances from roosts is important for land managers in the Perth-Peel region, given that all three species have been recorded roosting and foraging in Perth-Peel, and given it is a major foraging region for Carnaby's cockatoos.

¹² Le Roux, C. 2017. Nocturnal roost tree, roost site and landscape characteristics of Carnaby's Black-Cockatoo (*Calyptorhynchus latirostris*) on the Swan Coastal Plain.

¹³ Here, 'resident' at a roost means returning to the same roost on consecutive nights. When black cockatoos are not 'resident' at a roost, they may use other roosts in the same area or undertake seasonal shifts, likely in response to local depletion of food resources.

Cultural significance of black cockatoos

Background

To gather perspectives of Noongar people from the Perth-Peel region on the cultural significance of black cockatoos, the *Keep Carnaby's Flying – Ngoolarks Forever* project held cultural engagement workshops in 2022–23 with Elders and senior Noongar people from the Perth-Peel region. Workshop participants included delegates from Ngangk Yira Institute for Change Elders Council, based at Murdoch University, as well as Winjan Aboriginal Corporation and other Elders.

Themes that emerged in the workshops are presented here to facilitate local government cultural engagement for black cockatoo conservation. Quotes from participants are presented in their own words, as collated by the project's Cultural Engagement Lead Barb Hostalek with support from the Ngangk Yira Institute for Change. It is acknowledged that there is diversity of culture, knowledge and perspectives within Noongar Nations.

Themes from the workshops

1. Desire for Noongar people to work with local governments on black cockatoo conservation

The Carnaby's project is important. Winjan Aboriginal Corporation's involvement in supporting the project is very important, and a serious matter for our community. We grew-up with the birdlife, the birds, and once you see them go, you miss it. And this project is a collaboration involving revegetation, put in...water stations, looking at important revegetation opportunities, working across the metro area on Noongar Country.

We all, as a collective and full community, need to take full responsibility. To us as Noongar People, this is a bird that is part of life. And seeing the destruction of that as part of clearing of land is a concern. Being part of a community [we] take responsibility through education awareness with the shires we will be working with, and the wider community.

Collaborating with councils is a model to change the way things are done. It's an opportunity to work together to save the Ngoolarks.

Brett Hill¹⁴

We are all responsible for boodjar – for planet earth, sky land and water [including] the protection of these birds.

Jade Maddox¹⁵

We all want to be part of the solution.

Millie Penny¹⁶

2. Cultural significance

Families had them as a totem – they were messengers of the changing of the seasons and the coming of the rains.

Jade Maddox¹⁷

When you first hear them, it's around Easter time. It used to tell me, the salmon are coming around to where I am in Mandurah. They tell me that.

Franklyn Nannup¹⁸

You know, you know years ago, years ago, I was told that when black cockatoos flew south they brought rains, and when they moved went flew back north they took the rain away.

Franklyn Nannup¹⁹

The "tail of two cockies": Karrak the red-tailed black cockatoo acquired its red tail markings on the tail from Ngo-lak, the white tailed Carnaby's cockatoo. Mulal the swamp hen was feeding...on a sedge, the roots of which ooze red sap, and he cut a reed and struck Ngo-lak across his back. When Ngo-lak spread his tail to defend his back, Mulal threw lumps of red sap at his tail. Ngo-lak...turned into Karrak, the red-tailed black cockatoo.

Derek Nannup²⁰

[The] spirituality of the bird has been lost and we need to revive that...who are sharing these stories.

Marie Taylor²¹

¹⁴ Noongar Elder and Winjan Board Director, Elders Yarning Circle, 2023

¹⁵ Senior Bindjareb woman, Elders Yarning Circle 2023

¹⁶ Noongar Elder and Middar Yorga Dance Troupe

¹⁷ Senior Bindjareb woman, Elders Yarning Circle 2023

¹⁸ Noongar Elder and Winjan Aboriginal Corporation Director, Elders Yarning Circle, 2023

¹⁹ Noongar Elder and Winjan Aboriginal Corporation Director, Elders Yarning Circle, 2023

²⁰ Senior Noongar man (DBCA Education Officer), Elders Yarning Circle, 2023

²¹ Whadjak/Ballardong Elder and resident of City of Kwinana, Elders Yarning Circle, 2023



3. Concern about black cockatoo declines and the importance of protecting boodjar (country)

Noongar Country is kwobidak boodjar (beautiful country). It is an internationally-recognised Global Biodiversity Hotspot, and the only home of the Ngoolark (Carnaby's cockatoos).

Barb Hostalek²²

The Black cockatoo, one with a red tail, one with a white tail. Seeing them fly around is a normal part of life growing up as a kid and now as an adult. Their welfare is a grave concern, their lifestyle being impacted on by the destruction of the land whether it's through development for housing, suburbia. The Southwest is a precious place on the planet, with animals and plant life and we as Noongar people have been here 50,000 years.

George Walley²³

Everything is interconnected, the birds are dying, the food is dying, their waters are going...land is flattened without a care. It all interconnects. We used to see mobs of them...It's very sad.

Jade Maddox²⁴

Once they're gone, you can't get them back.

Franklyn Nannup²⁵

4. Solutions

[At the workshops] Elders expressed support for saving black cockatoos by protecting remaining bushland, and planting more trees as homes and food sources. They expressed concern that land clearing should not include ongoing clearing of places for Ngoolarks to live.

Barb Hostalek²⁶

How do we get [land-owners] to appreciate a native garden.

Derek Nannup²⁷

I've got the yard ready to go [to plant food for Ngoolarks]. The council should step in...if you're willing to put in a native garden...give us the plants we want...there's bush medicine, there's bush tucker.

Chelsey Thomson²⁸

Conversation between Brett Hill²⁹ and Franklyn Nannup³⁰

Brett Hill: *The challenge to all families, households... We all go camping, go bush... How do we first connect, what do we hear, if we listen? And if some of that natural environment is gone...?*

Franklyn Nannup: *[then] you hear nothing.*

So the challenge is to look at all development, [even at] schools, home, before you cut down a tree. Think about the significance of that tree...to the Carnaby's, to our birds. If they're gone, it's all gone...the challenge is to plant a tree.

I went out, me and my wife was out there with the Activate the Wheatbelt [community revegetation] mob, and then in one week we planted over 200,000 trees.

Mort Hansen³¹

22 Aboriginal woman and Cultural Lead, *Keep Carnaby's Flying – Ngoolarks Forever*, 2023

23 Bindjareb Noongar Elder and Winjan Board Director, Elders Yarning Circle 2023

24 Senior Bindjareb woman, Elders Yarning Circle 2023

25 Noongar Elder and Winjan Aboriginal Corporation Director, Elders Yarning Circle, 2023

26 Aboriginal woman and Cultural Lead, *Keep Carnaby's Flying – Ngoolarks Forever*, 2023


27 Senior Noongar man and DBCA Officer, Elders Yarning Circle, 2023

28 Whadjak woman and resident of City of Kwinana, Elders Yarning Circle, 2023

29 Noongar Elder and Winjan Board Director, Elders Yarning Circle, 2023

30 Noongar Elder and Winjan Aboriginal Corporation Director, Elders Yarning Circle, 2023

31 Noongar Elder, Elders Yarning Circle 2023



Importance of the City of Kwinana for black cockatoos

The City of Kwinana is one of the most important areas for black cockatoos in the Perth-Peel region. The City hosts both Carnaby's cockatoos and forest red-tailed black cockatoos³², with occupancy across its many roost sites placing it amongst the top LGAs in Perth-Peel for numbers of black cockatoos. This is due in large part to the presence of important black cockatoo foraging habitat in the City's reserves and green spaces.

The City of Kwinana has retained important roosting habitat to support large flocks of black cockatoos, and key food resources nearby to ensure that the roosts remain viable. **Figure 2** shows the known³³ black cockatoo roost sites in the City of Kwinana, as well as known roosts just outside the City's boundary, as some of the City of Kwinana's foraging flocks may roost just outside the City boundary at night.




Figure 2. Known black cockatoo roost sites within the City of Kwinana

Of the 15 known night roosts, all 15 have recorded use by white-tailed black cockatoos (either Carnaby's cockatoos or Baudin's cockatoos), and three have recorded use by forest red-tailed black cockatoos (some are used by more than one species). Roost locations are identified through BirdLife Australia's annual Great Cocky Count (early April) and Murdoch University flock tracking data; however not all flocks have been tracked, and while most roost sites in urban areas of Perth and Peel have been identified, it is possible that some have not, or that new sites may be used in the future.

Note: roosts located just outside the City are included in the map because some of the black cockatoos that forage within the City of Kwinana may use a night roost just outside the City.

Source: Data from BirdLife Australia's annual Great Cocky Count and flock tracking data from Murdoch University's Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).

³² The southwest part of the City of Kwinana is also within the likely distribution range for Baudin's cockatoos.

³³ 'Known' roosts are roosts identified by Murdoch University's tracking data, or roosts confirmed to be in use during BirdLife Australia's annual Great Cocky Count, which occurs on a single night in April each year. There may be additional roosts which are used by flocks at times of year not captured by the tracking and monitoring data.





The importance of the City of Kwinana for black cockatoos presents the City with opportunities and responsibilities. Local governments have considerable capacity to ensure that their communities continue to enjoy wild black cockatoos in their local area, given the powers of local government as land managers and environmental decision-makers at the local level. As black cockatoos continue to decline across their ranges, it becomes more critical to support the flocks that roost and forage in the Perth-Peel region.

The plight of black cockatoos is of increasing public interest, with community expectations that local governments will make decisions and take actions that align with scientific knowledge about the threats to black cockatoos and about what is required to retain them in the region and in specific local areas. This Black Cockatoo Conservation Action Plan will support the City of Kwinana to take an evidence-informed approach to the sustainable management of its black cockatoo flocks.

Legislative requirements and policy implementation

This Plan will also support the City of Kwinana to meet its legislative requirements regarding black cockatoos, and to implement the environmental and natural resource management policies of state and Commonwealth government.

Opportunities to leverage black cockatoo actions to meet Native Vegetation Policy outcomes

The Native Vegetation Policy (NVP) for Western Australia includes the policy outcome of 'net gain of native vegetation' and the strategic action of 'protection and enhancement of urban forests and bushland'. This Black Cockatoo Conservation Action Plan can assist the City of Kwinana to implement the NVP in a way that directly supports black cockatoos. For example, this Plan can ensure that the City's actions towards 'gain of native vegetation' and 'protection and enhancement of urban forests and bushland' (from the NVP) involve planting and protecting the specific native trees that are key foods for black cockatoos, and that these actions occur in locations that will be of value to black cockatoos, based on City-specific data.

Alignment with the City's existing strategies, policies and plans

Some information in this Plan will be familiar to City of Kwinana officers and elected representatives, and may already be incorporated in the City's strategies, policies, plans and guidelines. This Plan also includes new City-specific data and information from analyses undertaken as part of developing this Plan, specific to the City of Kwinana. This information is provided to inform and, where required, update existing strategies, policies and plans.

SMART goals: The actions in this plan will also allow the City of Kwinana to develop and implement SMART goals specific to its requirements.

The City of Kwinana's black cockatoos are a significant environmental and cultural asset for the City. Undertaking the actions in this Plan to protect and manage the City's flocks of black cockatoos will help the City of Kwinana to be a truly sustainable City for black cockatoos, while simultaneously enhancing urban forest, native vegetation, urban cooling, climate mitigation and local biodiversity.

Threat 1. Loss of habitat

Loss of habitat is the main direct cause of the declines of black cockatoos in southwest Western Australia. Clearing of native vegetation has stripped away too much of the birds' foraging, roosting and breeding habitat, and has fragmented and degraded much of what is left.

The formal conservation status for Carnaby's cockatoos, Baudin's cockatoos and forest red-tailed black cockatoos indicates that if net habitat loss continues, all three species will become extinct (within decades, for at least the first two species). Despite this, clearing of their remaining habitat continues, perpetuating their declines.

In Perth and Peel, local governments have unique powers and capacity to help black cockatoos in the region, due to their roles and responsibilities as land managers and decision-makers at the local level, and their roles in the sustainable management of biodiversity and native vegetation in their local area. To help the City of Kwinana to safeguard its black cockatoo flocks, this part of the Conservation Action Plan outlines the key habitat-related threats that are relevant for the area, and specific actions for the City to take to address these threats.



1.1 Lack of food resources

Lack of food is an identified threat to black cockatoos.³⁴ In Perth and Peel, flocks are under particular pressure.³⁵ Their main food resources, including jarrah, marri, banksia and hakea, have been extensively cleared. The threat of food scarcity in the Perth-Peel region has been exacerbated further by the clearing of the majority of the pine plantations, which had become a critical resource for Carnaby's cockatoos. In some areas of Perth, a lack of food has meant that black cockatoo flocks are already gone.

Areas without enough food resources lose their black cockatoos

Two Perth suburbs are shown in **Figure 3** and **Figure 4**, with black cockatoo flock movement data for each area. In **Figure 3**, development within a suburb has been poorly designed and managed with respect to retention of green spaces and native vegetation; without enough food, black cockatoos have been lost from the area. In **Figure 4**, retention of native vegetation, green spaces and 'vegetation corridors' (vegetation patches within several hundred metres of each other, which act as stepping stones between larger areas of foraging habitat), and larger suburban blocks with gardens, have allowed black cockatoos to continue to forage. This area has kept its black cockatoos.

³⁴ The national recovery plans for southwest Western Australia's black cockatoos identify the need to plant more foraging habitat, if declines are to be reversed.

³⁵ Williams et al. (2017); BirdLife Australia (2023).



Figure 3. GPS data showing foraging movements for a tagged Carnaby's cockatoo and its flock

The birds use the retained green spaces and vegetation patches for foraging. Note also the use of a horizontal belt of retained tuart trees running through adjacent residential plots, which the flock used as corridor habitat to allow access to foraging habitat on the western side of the suburb. The densely built-up areas, where vegetation has been removed and not replanted, are devoid of flock movement. Black cockatoos have been lost from these areas.

Source: Data from Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).



Figure 4. GPS data showing foraging movements for a tagged Carnaby's cockatoo and its flock

This figure shows a continued high presence of black cockatoos in the suburban areas with larger blocks and retained vegetation (east side of map). In contrast, the newer housing development in the centre of the map has been poorly planned and designed with respect to native vegetation (as well as urban canopy, urban cooling and public open spaces), and involved the removal of all vegetation. This area provides no foraging or movement opportunities for black cockatoos, and is largely devoid of black cockatoo activity and presence.

Source: data from Murdoch University Black Cockatoo Conservation Management Project). Figure produced using Google Earth Pro (<https://earth.google.com>).



Black cockatoos need food in close proximity to their roosts

Flock tracking data show that outside of breeding migrations, black cockatoo flocks in the Perth-Peel region typically spend weeks to months roosting in a particular area, during which time the birds forage in close proximity to that roost.³⁶ In general, the *maximum* daily distance that flocks will forage from their resident roost is around 6km³⁷. This highlights the need for local governments to ensure that black cockatoos have enough food resources in close proximity to their roosts. If there is ongoing net loss of foraging habitat within 6km of a roost, eventually flocks will be lost from that area.

Black cockatoos need food security during times of low food yield

It is critical to recognise that much of the foraging habitat in an area may not produce food for black cockatoos at certain times of year, due to seasonal restrictions in fruiting and flowering. Patches that contain a range of food species with different fruiting months will mitigate this risk, however there can also be large variations in food production between years. In some years, banksia woodland on the Swan Coastal Plain produces only half the fruit yield of other years³⁸, greatly reducing the total food available for black cockatoos, and individual marri trees only have a high fruit yield every 3–5 years³⁹. Flocks need enough foraging

habitat close to their roosts (within 6km) to ensure they still have food available during months and years of lower food productivity.

If seasonal and interannual differences in food availability are not planned for and managed effectively by natural resource managers, then during low-yield periods, black cockatoos are at risk of depleting all available food resources within daily foraging range of their roosts.

The value of small patches is often ignored

In the Perth-Peel region, where larger areas of bushland are scarce, black cockatoos can only survive by making extensive use of small patches of food resources. Murdoch University's flock tracking research shows that no patch is too small to be used as a food resource by black cockatoos⁴⁰. Small patches of foraging habitat and single food trees include those in parks and other green spaces, around ovals, along road verges and in private backyards. These small patches are often undervalued and at risk of being cleared; which can lead to large cumulative losses of food resources. All are important, to retain black cockatoos in urban areas.

Quality of foraging habitat can be misjudged

Quality of native vegetation as a food resource for black cockatoos should not be assessed using general 'vegetation quality' ratings. Food trees remain valuable for black cockatoos even if the understorey is 'degraded'. Likewise, Murdoch University flock tracking research shows that in urban areas, black cockatoos often feed on small patches of banksia that have relatively 'sparse' canopy cover. These vegetation conditions, which may be rated 'low' quality in other contexts, often represent high value food resources in the Perth-Peel region. Failure to recognise this can lead to the clearing of important food resources which should have been either retained, or replaced – by planting new food sources nearby before clearing occurs – to avoid net loss for flocks in the local area.

Plant diseases, fire and climate-related risks

Plant diseases such as dieback and marri canker, and risks from fire and climate impacts all threaten the viability of black cockatoo foraging habitat. The risk increases when areas of habitat become small and more exposed to external risks; as is the case in the Perth-Peel region. Plant diseases, fire and climate changes threaten not only foraging habitat but also roosting and breeding habitat, and as such, each of these threats is covered separately, later in this Plan.

36 Murdoch University's flock tracking data and BirdLife Australia's Great Cockey Count data show that flocks display seasonal residence patterns, with the same roosts used year after year. Flocks departing the Town of Cambridge for breeding areas return to these same roosts with their newly fledged young beginning in late summer, in anticipation that the same patches of food resources will be there to support them.

37 Murdoch University data for the Perth-Peel region and Swan Coastal Plain show that when a Carnaby's cockatoo flock is 'resident' at a roost (returning to the same roost each night), the flock forages at a mean distance of 4.26 +/- 2.24km from the roost (Murdoch University unpubl. data 2023). These findings support previously reported daily maximum foraging distances of around 6km from roosts (e.g. Kabat et al. 2012; Rycken et al. 2021). State and Commonwealth government conservation agencies including DBCA and DCCEEW also use this figure of <6km as a maximum daily foraging distance from roosts for black cockatoo flocks during the non-breeding season; the time when black cockatoos on the Swan Coastal Plain need to find enough food to build their energy reserves for the next breeding season.

38 See Appendix A of Williams et al. (2017) for discussion of recurring low-yield years for banksia.

39 See, for example, Robinson 1960; Mawson 1995; Johnstone & Kirkby 1999.

40 Riley et al. (2023) unpubl. data

Cumulative impacts

The numerous small and larger clearing actions in the Perth-Peel region, on both public and private land, represent ‘death by a thousand cuts’ for black cockatoos. Many small net losses of food will eventually mean an area becomes unviable for black cockatoo flocks. Yet the impacts of a clearing action are still often assessed from the point of view of the ‘individual’ impact, without appropriate consideration of cumulative impacts. This occurs despite regulatory requirements to consider cumulative impacts.

Nor are the impacts of clearing a patch of foraging habitat typically considered at the level of the individual flock. It is important not only that actions are undertaken to mitigate the loss of food from clearing, but that these actions are implemented close to the cleared site, to help support the local flocks. The species declines of Western Australia’s black cockatoos are being led by declines of individual flocks, and managing natural resources to ensure no net loss of food for individual flocks will help to halt these declines.

Inadequacy of planning laws

Perth has the lowest retention of urban canopy of any capital city in Australia, and government has recognised that planning laws have been inadequate to enable tree retention⁴¹. The WALGA recognises the need for net gain of canopy and particularly native vegetation in urban areas. In 2023, the WALGA asked state government to increase funding for the Urban Forest Grant Program, to help local governments plant more trees in their local areas. This aligns strongly with the need to plant more food trees for black cockatoos.

⁴¹ Minister for Planning, Hon Rita Saffioti MP (2023), as quoted in www.ac.net.au/news/2023-03-11/perth-housing-developments-that-could-improve-tree-canopy/102060214



Actions »

To address the threat from lack of food, there are key actions that the City of Kwinana can take. Across Perth and Peel, governments must halt and reverse the ongoing ‘net loss’ of food resources. In the City of Kwinana, this means *retaining and improving* the remaining food resources, as well as creating additional food for black cockatoos.

These actions are described over the next several pages, supported by LGA-specific data and maps to enable implementation. These actions to halt and reverse the loss of food resources are the most important direct actions that the City of Kwinana can take to safeguard its black cockatoos.

Retain and improve existing food resources

- » Retain black cockatoo foraging habitat on council managed and regulated land.
- » While creating new habitat is critical, retention of existing habitat is the highest priority, because:
 1. When new food resources are planted to offset the clearing of existing food resources, there is a ‘food gap’ of at least 5–10 years before the new plants begin providing food. Given that food insufficiency is already causing black cockatoo declines, any additional food shortage will increase the risks to flocks in the local area, and the species⁴² overall.
 2. Revegetated areas of foraging habitat such as banksia woodland are often of lower foraging value than the original cleared vegetation, leading to net loss of quantity and quality of food for flocks in the local area⁴³. Retention of existing resources avoids this problem.
 3. Success of revegetation is not guaranteed, and all efforts require long-term management. Retention of existing habitat avoids these issues and risks.⁴⁴
- » Improve the condition and quality of existing foraging habitat, through infill planting, weed and disease control (dieback, marri canker), and fire risk management.
- » Recognise that the quality of foraging habitat for black cockatoos should not be assessed by general ‘vegetation quality’ ratings. Food trees remain valuable for black cockatoos even if the understorey is ‘degraded’; in urban areas, black cockatoos often forage on small patches of banksia that have relatively ‘sparse’ canopy cover. These are still high value food resources in the Perth-Peel region.
- » Retain small remnants of native vegetation. Even small patches may contain a diversity of food trees that will provide food at different times of year. In Perth and Peel, remnant patches of food are valuable however small.
- » Retain individual large food trees (e.g. jarrah, marri). Their value as food (and roosts) is difficult to replace.

42 To avoid impact on flocks from clearing of food resources, some clearing proponents proactively plant new foraging habitat in the area of the proposed clearing action (within 6km), several years before clearing is proposed to occur. This is often an appropriate mitigation, as it aims to safeguard flocks from additional food shortages that could have arisen from the clearing action. Six kilometres aligns with the approximate upper limit of a flock’s daily foraging distance from their roost (see details in earlier sections).

This mitigation also aligns with the EPA’s technical advice regarding Carnaby’s cockatoo foraging habitat in Perth and Peel (EPA 2019). The EPA advice states that any clearing should be offset by undertaking revegetation, or rehabilitation of degraded areas, within “close proximity of the impacted area”, to provide alternative food resources for the flocks affected by the action. The EPA advice acknowledges that such actions will take time to provide new food resources; highlighting the need to commence planting 5–8 years before clearing, to allow time for replacement food to become available.

43 Although this risk can be mitigated by replanting over a larger area than the original cleared area and planting for species diversity, it is challenging to find enough space for additional revegetation in urban areas, particularly given that flocks affected by a clearing action need replacement habitat within their daily foraging range (4–6km from their roosts).

44 Occasionally there will be no option but to clear food resources. In such cases, local governments should require, as a condition of clearing, that the proponent ensures no net loss (and preferably net gain) of foraging resources for flocks in the local area. For simplicity, ‘local area’ is within 6km of the clearing action (this aligns with research regarding daily foraging distances for flocks on the Swan Coastal Plain). This approach was adopted by the City of Canning in a recent clearing referral. The City submitted a referral to clear 2.4ha of black cockatoo foraging habitat, but ensured it had already created an ‘advanced offset’. Prior to submitting the referral, the City had created 6ha of new foraging habitat on degraded land at a Bush Forever site within 6km of the proposed clearing area. The City recognised that three points are essential to mitigate clearing of black cockatoo food resources:

- new foraging habitat must be created to offset what is lost (rather than only acquiring existing bushland, which perpetuates net loss).
- the new habitat must be created within the daily foraging range of affected flocks, to ensure no impact to those flocks.
- the new habitat should be established several years prior to clearing, so that it is mature enough to be providing food before any food is removed, thereby avoiding years of ‘net loss’ which could impact flocks.

In this example, the City recognised that if ongoing net habitat loss led to insufficient food for flocks within foraging range of their roosts, the roosts would become unviable and black cockatoos would be lost from the area. The City took effective action to prevent this.

- » Retain pine trees and Cape Lilac until alternative food sources become available. In the Perth-Peel region, where native food sources are scarce, pine is a high-energy food resource for Carnaby's cockatoos, and Cape Lilac is important for forest red-tailed black cockatoos⁴⁵.
- » Retain food resources on private land
 - Large losses of native vegetation and tree canopy are due to clearing on private land⁴⁶. Importantly, local government regulators and decision-makers have influence over much of this clearing; such as through rezoning decisions. Local governments can support black cockatoos by making decisions which retain local food resources.
 - At time of writing, many local governments in Perth are advocating for legislation to help them identify and protect 'important trees' on private property, for the trees' aesthetic value as well as for urban cooling and local biodiversity. Support for this legislation is support for black cockatoos.
- » Apply formal measures and mechanisms to enable the City of Kwinana to better retain black cockatoo food resources on government and private land, including land management agreements and strategies as required. This will support legislative responsibilities (e.g. BC Act, EPBC Act), and complement the City of Kwinana's roles, responsibilities and strategies in related areas (e.g. urban biodiversity; urban forest/canopy; urban cooling; climate mitigation).



- » Undertake community awareness-raising about the value and necessity of retaining food resources on government and private property, if suburbs are to retain their black cockatoos. Black cockatoos are charismatic flagship species for community engagement efforts.
 - Important community messages include that black cockatoos in the City of Kwinana make extensive use of smaller areas of green space and private gardens when feeding.

Pink-flowering marri: They have attractive flowers and help feed black cockatoos in urban landscapes.

⁴⁵ Taking pine as an example, this is important in areas where there may be food shortages, due to the high energy content of pine seed and the large number of cones per tree. The 'carrying capacity' of pine forest is 9x greater than banksia woodlands, i.e. for the same sized area of vegetation, pine supports 9x as many Carnaby's cockatoos as banksia woodland (Williams et al. 2017). However, native foods like banksia provide essential proteins and micronutrients, and tracking studies show that Carnaby's cockatoo flocks that feed in pine plantations still spend part of each day feeding on native foods (Murdoch University unpubl. data 2023).

⁴⁶ Western Australian Department of Parks and Wildlife, (2013). *Carnaby's cockatoo Recovery Plan*.

Create additional food resources

On the Swan Coastal Plain, where food is already scarce, black cockatoo declines will not be reversed without shifting to 'net gain' of food resources. This Plan provides information about *what* to plant, and *where* to plant, for the City of Kwinana's black cockatoo flocks.

What to plant

- » Plant species from the Proteaceae family, particularly banksia and hakea, to provide food in 5–10 years and address current food shortages. Plant seedlings rather than seeds.
 - » Plant marri and jarrah, as established trees where possible, for longer-term food resources (and roosts). These two trees are critical for all three black cockatoo species, as although Carnaby's cockatoos eat banksia and hakea as well as marri and jarrah, Baudin's cockatoos depend almost solely on marri as a native food source, and forest red-tailed black cockatoos depend on marri and jarrah.
 - » In urban areas where food resources are scarce, black cockatoos also benefit from introduced pine and macadamia (Carnaby's cockatoos) and Cape Lilac (forest red-tailed black cockatoos). Plant macadamia for a fast-growing food resource⁴⁷.
- » Ensure all planting and plant maintenance are planned and resourced for optimal plant survival.
 - » Plan for infill planting to replace losses.
 - » Plant a variety of species (**Table 1**).
 - Different plants provide food for black cockatoos at different times of year. Access to a variety of food species will support flocks in the City of Kwinana during times of year when some plants are not providing food.
 - Different plants provide different important nutrients⁴⁸.
 - » Plant many of the same species (with the same fruiting times) in the same area, so that sufficient food becomes available at the same time of year to feed an entire flock.
 - » Where possible, avoid herbicides and pesticides on black cockatoo food plants, particularly when the plants are producing food resources (i.e. flowers, fruits).
 - » Assess whether dieback is present at the proposed planting site. This will influence what to plant, given that some plants, particularly Proteaceae, are susceptible to this disease⁴⁹.
 - » Plant banksia species in well-drained areas to minimise losses from dieback.



47 Macadamia is recommended as a fast-growing, high-energy food source for Carnaby's cockatoos, given that there is not enough native foraging habitat left on the Swan Coastal Plain to halt their declines. Other common nut trees in the Perth-Peel region are not useful to plant for black cockatoos, because they have nuts which can be eaten by smaller cockatoo species such as corellas, which outnumber black cockatoos and may take the majority of the food. Macadamia nuts have hard shells which only black cockatoos can break.

48 As an example, Murdoch University's tracking of Carnaby's cockatoo flocks in the Perth-Peel region has found that the birds may feed heavily on plantation pines at times of year when cones are available, but will always also spend some time each day foraging on banksia and other native species. While pine is a high-energy food for Carnaby's cockatoos, banksia provides them with important proteins and micro-nutrients.

49 The Dieback Working Group has resources and publications available on their website regarding the relative resilience of various local native plants to dieback, as well as about how best to manage dieback in bushland, and other useful information: www.dwg.org.au

Table 1. High priority food plants to plant for black cockatoos in the City of Kwinana^{a,b}

Tree type	Species name	Height (m)	Flower colour; flowering season	Importance for which black cockatoo species
Banksias	Slender banksia (<i>Banksia attenuata</i>)	Medium tree; 1–10m	Yellow; Oct–Feb	• Carnaby's cockatoo
	Swamp banksia (<i>Banksia littoralis</i>)	Medium shrub; 1–2m	Yellow; Mar–July	• Carnaby's cockatoo
	Firewood banksia (<i>Banksia menziesii</i>)	Medium tree; 8–10m	Red/yellow or pink/yellow; Feb–Oct	• Carnaby's cockatoo
	Acorn banksia (<i>Banksia prionotes</i>)	Medium tree; 8–10m	Bright orange; Feb–Aug	• Carnaby's cockatoo
	Oak-leaved banksia (<i>Banksia quercifolia</i>)	Large shrub; 1.5–3m	Orange/yellow; Mar–Nov	• Carnaby's cockatoo
	Showy banksia (<i>Banksia speciosa</i>)	Small to medium tree; 1–6m	Cream or lemon yellow; year-round	• Carnaby's cockatoo
	Parrot bush (<i>Banksia sessilis</i>)	Large shrub; 0.5–5m	Yellow-cream; Apr–Nov	• Carnaby's cockatoo
Hakeas	Two-leaved hakea (<i>Hakea trifurcata</i>)	Large shrub; 1.5–3m	Pink or white-cream; Apr–Oct	• Carnaby's cockatoo
	Wavy-leaved hakea (<i>Hakea undulata</i>)	Medium shrub; 1–2m	White; July–Oct	• Carnaby's cockatoo
Larger trees	Marri (<i>Cormbica calophylla</i>)	Large tree; 40m	Pink or white; Dec–May	• FRTB cockatoo • Carnaby's cockatoo
	Jarrah (<i>Eucalyptus marginata</i>)	Large tree; 40m	White-cream or pink; June–Jan	• FRTB cockatoo • Carnaby's cockatoo
	Coastal blackbutt or Pricklybark (<i>Eucalyptus todtiana</i>)	Medium tree; up to 15m	White, Jan–April	• Carnaby's cockatoo
	Macadamia (<i>Macadamia integrifolia</i>)	Medium tree; 2–12m depending on species	Varies by species (purple, pink or white); flowering and fruiting times vary by species	• Carnaby's cockatoo
	Pine ^c (<i>Pinus</i> sp.)	Large tree; varies by species	Varies by species; flowering and fruiting times vary by species	• Carnaby's cockatoo

Note: (a) For additional suitable black cockatoo food species and information on plant selection and the design of plantings, see resources on the website of BirdLife Australia.

(b) Re polyphagous shot-hole borer (PSHB): As of 2024, DPIRD's advice (pers. comm. BirdLife Australia 2024) is: "the only trees DPIRD recommends avoiding [planting] are the Preferred Reproductive Host trees for PSHB". It is important to note that the term 'Preferred Reproductive Host trees' refers only to **the first 10 trees listed** in the following document (current September 2024): www.agric.wa.gov.au/sites/gateway/files/PSHB-WA-Host-List.pdf. The 10 trees on the 'Preferred Reproductive Host trees' list: Box elder maple; Mirror bush; Poinciana; Coral tree; Moreton Bay fig; Port Jackson fig; White mulberry; Black mulberry; London plane tree; Robinia. None of the black cockatoo food plants in Table 1 is a 'Preferred Reproductive Host' tree, and therefore all are acceptable to plant, as per this 2024 DPIRD advice. Contact DPIRD for latest advice.

(c) *P. pinaster* is the preference; a Carnaby's cockatoo can meet daily energy requirements from 11 cones of *P. pinaster*, or 18 cones of *P. radiata* (Stock et al. 2013). They will feed on all pine species, including native pine.



Remember to water regularly

- » Although native trees are waterwise plants, all young trees need regular watering in their first 2–3 years, while they establish.
- » Without suitable watering regimes, young tree losses can be catastrophic.
- » During warmer weather, young trees may need watering 1–3 times per week, depending on species.
- » If outsourcing plant management, ensure responsibility for watering is clear, and includes infill planting to replace losses.
- » Even established trees may need regular deep-watering in summer; particularly during heatwaves. Wetting agents may be useful; follow best practice for species and conditions.
- » Contact local NGOs (e.g. Urban Bushland Council) and Friends groups for support with watering.

**BLACK
COCKATOO
CONSERVATION**



ACTIONS

Where to plant

Council-managed green spaces

- » Provide additional black cockatoo food resources in council-managed green spaces. Extend native vegetation patches; enhance with infill planting; and create new patches. Creating new bushland in POS is a WALGA priority⁵⁰.
- » New habitat is useful even as small areas. Food patches of 0.1–0.5ha are used by black cockatoos, as are single trees.
- » Consider planting patches of banksia and hakea, and advanced tree planting of marri, jarrah and macadamia, around parks and open spaces.


To support these actions, the locations of City-managed green spaces in relation to known roosts are shown in **Figure 5**. Planting food within daily foraging distance of roosts (not more than 4–6km) will help the City's flocks.

Figure 5. City of Kwinana showing council-managed green spaces in relation to known night roosts for black cockatoos

Green spaces near roosts are of value to investigate for revegetation, infill planting and advanced tree planting with key food species. Roosts just outside the City are included in the map, as black cockatoos that forage within the City may use roosts just outside.

Source: Green space data obtained from spatial layers provided by the City ('All Reserves within Kwinana.shp' & 'City of Kwinana Managed Reserves_Local Biodiversity Strategy.shp'; obtained 30/04/2024). Black cockatoo data from BirdLife Australia and Murdoch University. Figure produced using Google Earth Pro (<https://earth.google.com>).

Note: High resolution maps are available. See page 5 of this document for more information.



Bush Forever areas and DBCA land

Bush Forever areas and DBCA land contain remnant native vegetation and food resources that are critical for retaining black cockatoos in local areas.

- » Working with DBCA as required, undertake revegetation activities and infill planting with black cockatoo food species in Bush Forever areas and DBCA land.
- » Small patches are important.
- » Improve the condition and connectivity of existing habitat.

To support these actions, the locations of Bush Forever areas and DBCA land in the City of Kwinana in relation to known black cockatoo roost sites are shown in **Figure 6** and **Figure 7**. Provision of additional food within daily foraging distance of roosts (not more than 4–6km) is a critical action to support the City’s black cockatoo flocks.

Figure 6. City of Kwinana: Bush Forever areas in relation to known roost sites for black cockatoo flocks, showing areas of value for the City to investigate for revegetation activities and infill planting with black cockatoo food species

Roosts located just outside the City are included in the map, because the black cockatoos that forage within the City may use night roosts just outside the City.

Source: Bush Forever areas were obtained from the Department of Planning, Lands and Heritage data layer ‘Region Scheme – Special Areas (DPLH-022)’ (catalogue.data.wa.gov.au/dataset/region-scheme-special-areas-dop-073; accessed 07/02/2024). Black cockatoo data are from BirdLife Australia and Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).

Note: High resolution maps are available. See page 5 of this document for more information. 





Figure 7. City of Kwinana: DBCA reserves and legislated lands in relation to known roost sites for black cockatoo flocks*

Showing areas of value to investigate for local government to work with DBCA on revegetation activities and infill planting with black cockatoo food species. Roosts located just outside the City are included in the map, because the black cockatoos that forage within the City may use night roosts just outside the City.

*Note: * While most night roost sites in the City of Kwinana are likely to have been located, it is possible that some roosts remain unidentified, or that new sites are used in the future.*

Source: DBCA lands were obtained from the Department of Biodiversity, Conservation and Attractions data layers 'DBCA - Legislated Lands and Waters (DBCA-011)' (catalogue.data.wa.gov.au/dataset/dbca-legislated-lands-and-waters; accessed 06/02/2024) and 'Regional Parks (DBCA-026)' (catalogue.data.wa.gov.au/dataset/regional-parks; accessed 06/02/2024). Black cockatoo data are from BirdLife Australia and Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).

Note: High resolution maps are available. See page 5 of this document for more information.



BLACK COCKATOO CONSERVATION



Street trees and road verges

In addition to planting food trees in the council-managed green spaces identified in **Figure 5**, the City of Kwinana can support local black cockatoo flocks by planting food tree species as street trees and on road verges. These are valuable areas for food resources for black cockatoos in urban landscapes.

Safe set-backs from roads

All planting of black cockatoo food trees near roads should include appropriate set-back distances, so birds can access the trees safely:

- » In quiet suburban streets (speed limit <50km/hr), plant food trees at least 3–5m back from road edges. Planting along a single side of the road will reduce the risk of birds flying across the road when moving between trees.
- » For busy roads (speed limit >50km/hr), plant food trees at least 10m from road edges.
- » Maintain existing large food trees (jarrah, marri) on verges by pruning low branches.
- » Never plant black cockatoo food plants on median strips; this is a death trap for the birds.

Connectivity

Create vegetation corridors to help flocks access food. Black cockatoos prefer to make short 'step' flights between neighbouring patches of habitat, when moving through fragmented urban landscapes (see vegetation corridors section of this Plan). If visual inspection of the maps in this Plan reveals areas between roosts and existing foraging habitat where native vegetation is low or absent, consider creating a small patch of foraging vegetation there to act as a stepping stone.




Figure 8. Schools and their ovals often provide valuable foraging and roosting habitat for black cockatoos

In this example, Murdoch University flock tracking data reveals two GPS-tagged black cockatoos and their flocks making use of remnant bushland and individual food trees on the grounds of a Perth school for feeding, highlighting the value of planting black cockatoo food resources on school grounds.

Note: The two different colours indicate foraging movements for the two tagged birds, each in a separate flock, during the limited time that the tags remained operational.

Source: GPS tracking data are from Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).



School land

Plant additional food resources on school land. GPS and Argos satellite telemetry tracking of black cockatoo flocks by Murdoch University has revealed that remnant native vegetation on school land is an important food resource for these flocks. **Figure 8** is provided as an example.

- » Engage with schools and state government to undertake revegetation activities and infill planting on school land, using black cockatoo food species.
- » Even small patches are valuable.
- » City of Kwinana community engagement teams can use black cockatoos as flagship species, and hooks such as ‘create a Carnaby’s café’⁵¹ to activate school communities.

51 For planting a Carnaby’s café, see www.murdoch.edu.au/research/hbi/carnabys/how-you-can-help/help-by-planting-food



Priority areas for revegetation

To help prioritise where to plant additional food resources in the City of Kwinana, the following three maps show the 10 largest known black cockatoo roosts (by number of birds) for flocks that forage in the City of Kwinana, with a 2km buffer around each of these roosts (**Figure 9**).


Areas within these buffers that may be suitable for planting additional food are identified as 'priority areas'; to focus expenditure of limited resources in the short term. They should be considered as a guide only: while flocks prefer to forage close to roosts, they will forage up to around 4–6km from their roost, and all revegetation and restoration activities within the City of Kwinana will benefit local flocks. For the same reason, all foraging habitat that remains within the City of Kwinana is critical to retain.

Figure 9. Recommended priority areas for habitat retention and revegetation in the City of Kwinana

Continued next page

Priority areas were identified using 2km buffers (lime rings) around the largest known roosts (by number of birds) for flocks that forage in the City of Kwinana. No map is included for the northwest quadrant, because that quadrant is not within any 2km foraging buffers around the 10 largest known roosts. **Native revegetation areas:** include areas of bushland potentially suitable for infill planting with native tube stock. **Parkland revegetation areas:** include playgrounds, parks, ovals or other open spaces potentially suitable for parkland tree planting with either native or exotic tree species. **Mixed revegetation areas:** include a mixture of either remnant bushland/cultivated garden beds potentially suitable for native tube stock planting, and public open space potentially suitable for parkland tree planting with either native or exotic tree species. **Pine:** provides important foraging opportunities for Carnaby's cockatoos, therefore the locations of pine stands with 5+ trees are identified on the map (excluding stands on private property, as far as could be established).

Source: Black cockatoo data are from BirdLife Australia and Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).



**Note: While these priority areas have been provided to focus expenditure of limited resources, flocks will forage up to around 4–6km from their roosts, and all habitat within the City should be considered critical for the retention of flocks, alongside a policy of 'no net loss' of foraging habitat within the City's boundaries.*

Note: High resolution maps are available. See page 5 of this document for more information.




Figure 9. Recommended priority areas for habitat retention and revegetation in the City of Kwinana (cont.)

Priority areas were identified using 2km buffers (lime rings) around the largest known roosts (by number of birds) for flocks that forage in the City of Kwinana. **Native revegetation areas:** include areas of bushland potentially suitable for infill planting with native tube stock. **Parkland revegetation areas:** include playgrounds, parks, ovals or other open spaces potentially suitable for parkland tree planting with either native or exotic tree species. **Mixed revegetation areas:** include a mixture of either remnant bushland/cultivated garden beds potentially suitable for native tube stock planting, and public open space potentially suitable for parkland tree planting with either native or exotic tree species. **Pine:** provides important foraging opportunities for Carnaby's cockatoos, therefore the locations of pine stands with 5+ trees are identified on the map (excluding stands on private property, as far as could be established).

Note: While these priority areas have been provided to focus expenditure of limited resources, flocks will forage up to around 4–6km from their roosts, and all habitat within the City should be considered critical for the retention of flocks, alongside a policy of 'no net loss' of foraging habitat within the City's boundaries. Note 2: For the City's southeast quadrant specifically, not much city-managed green space exists within 2km of the largest roosts (i.e. the roosts with 2km buffer rings, within which we generally advise to prioritise revegetation). Therefore, to support black cockatoos using this quadrant, we recommend extending priority revegetation efforts to include city-managed green spaces that lie within 2km of **any roosts in this quadrant, not just the largest roosts.*

Source: Black cockatoo data are from BirdLife Australia and Murdoch University Black Cockatoo Conservation Management Project. Figure produced using Google Earth Pro (<https://earth.google.com>).

Note: High resolution maps are available. See page 5 of this document for more information.





Build black cockatoo-friendly actions into the City's strategies and policies

- » Building actions that support black cockatoo food resources into the City's strategies, policies and plans will enable a whole-of-City approach towards ensuring enough food for black cockatoo flocks in the City of Kwinana.
- » Key strategies, policies and plans to update to align with these actions may include those related to clearing proposals; urban forest/canopy; urban cooling and climate mitigation; biodiversity; street trees and road verge management; trees on private property; and public open space.
- » Set and maintain a formal target of **net gain** of foraging habitat in the City of Kwinana (public and private land), recognising that ongoing net loss will lead to loss of black cockatoos. Net gain will also benefit other threatened native birds and pollinator insects, and contribute to urban greening and canopy, beautification of streets and public open spaces, climate management and urban cooling.

Engage the community to plant black cockatoo food

- » Provide the community with information to help them protect black cockatoo food trees and give the birds more food, including planting suitable food in gardens and other private property (see BirdLife Australia's website for details).

Plant macadamia for a fast-growing food resource.

- » Use key dates such as World Bird Day (January), Clean-up Australia Day (March), International Day for Biodiversity (May) and National Tree Day (July) to create awareness and on-ground action. Dates in autumn and winter (planting season) can be leveraged to offer native plant and seedling giveaway and subsidy schemes to residents and Friends of Bushland groups, and to hold planting and rehabilitation days.
- » Organise and promote community planting days⁵², including in collaboration with land care NGOs⁵³, Friends groups and DBCA. Liaise with these organisations about events they are organising which the City could help promote and support.
- » Provide information about land care NGOs, Friends groups, BirdLife Australia and Kaarakin Black Cockatoo Conservation Centre, to activate the community to volunteer with these groups and assist in black cockatoo conservation actions.
- » Engage with businesses and utility providers to identify new areas for revegetation, such as on golf courses, school grounds and power easements.

Key points

- » Urban areas lose their black cockatoos when too much of their food is lost.
- » When resident at a roost, flocks need enough food as close as possible to the roost (within 4–6km).
- » Flocks of Carnaby's cockatoos, forest red-tailed black cockatoos and potentially Baudin's cockatoos roost and forage in the City of Kwinana.
- » Adopt a policy of 'net gain' of food resources. Without this, black cockatoo declines cannot be reversed.
- » Both habitat retention and creation of new habitat are important to safeguard the City's black cockatoos.
- » Engage land care NGOs and the community to help.
- » Use this Plan to update the City's strategies, plans and guidelines, to ensure they support the planting of black cockatoo food species.

⁵² In 2024, in conjunction with *Keep Carnaby's Flying – Ngoolarks Forever*, planting days for black cockatoo food were held in the City of Kwinana, in collaboration with natural resource management and land care NGOs. These days attracted dozens of community members, and several thousand seedlings were planted. This demonstrates the popularity of planting days.

⁵³ Recommended organisations include the partner organisations on the *Keep Carnaby's Flying – Ngoolarks Forever* project; see logos at the start of this document

1.2 Lack of roosts

Black cockatoo roost habitat is under threat across Perth and Peel, where most habitat suitable for roosting has been cleared. In some urban areas, no roost habitat remains.

What is suitable night roost habitat?

Black cockatoos roost communally at night in one or more tall, large trees; often the tallest trees in the area. Night roost sites can support more than one flock together, and different species may share a roost. Trees can be native or non-native; key requirements include that the trees must be tall (research shows a preference for >25m⁵⁴) with medium density foliage (around 50%). In areas where insufficient roost habitat remains at a single location, the flock may divide to roost in two or three locations that are within a few hundred metres of each other. Together, the locations are then considered a single roost.

During the time that a flock is resident in an area – for example, when a Carnaby's cockatoo flock is roosting and foraging in the City of Kwinana during the non-breeding season – the flock may use a few different night roosts within the area depending on food availability and weather conditions. Flock tracking data show flocks have preferred roosts, and revisit the same roost or network of roosts at a particular time of year each year, coinciding with seasonal patterns in food availability⁵⁵. This highlights the need for flocks to have several suitable roost locations in a particular area.



54 Le Roux, C. (2017). Nocturnal roost tree, roost site and landscape characteristics of Carnaby's Black-Cockatoo (*Calyptorhynchus latirostris*) on the Swan Coastal Plain.

55 Berry, P. F. (2008). Counts of Carnaby's Cockatoo (*Calyptorhynchus latirostris*) and records of flock composition at an overnight roosting site in metropolitan Perth. *Western Australian Naturalist*, 26(1), 1. Groom, C. (2015). *Roost site fidelity and resource use by Carnaby's cockatoo, Calyptorhynchus latirostris, on the Swan coastal plain, Western Australia* (Doctoral dissertation, University of Western Australia).

Roost viability

Night roosts must be supported by close proximity to enough food resources to feed the flock (see previous section) and by close proximity to safe drinking water (see next section). If either food or water becomes too scarce to support the flock, the roost becomes unviable.

Threats to roost habitat include:

- » Ongoing clearing of suitable tall trees on government and private land:
 - Few roost sites remain in each local government area in the Perth-Peel region.
 - Although just under two-thirds of known roosts in the Perth-Peel region are in parks and reserves, the remaining roosts are located on private land, highlighting the challenges of protecting roosts in urban areas.
 - Roosts are large trees. If removed, they cannot be replaced within decades.
- » Lack of enough food resources within daily foraging range of a roost (6km) to allow the roost to remain viable:
 - See previous section for the details of the research showing that black cockatoos need enough food within 6km of their roost to support all members of the flock.
 - See previous section for threats to food resources in the City of Kwinana, and actions to undertake.
- » Lack of reliable safe drinking water in close proximity to the roost (see next section).
- » Lack of succession planning for new roost habitat.
- » Disturbance at night.

Note: risks to roost viability from loud noises and bright lights at night require investigation. Researchers from Murdoch University's black cockatoo research team are looking at the impact of artificial light at night on roost quality.

Value of the City of Kwinana's roosts for black cockatoos

- » The City of Kwinana has night roosts for Carnaby's cockatoos, forest red-tailed black cockatoos, and potentially Baudin's cockatoos.
- » Some of the roosts comprise enough trees to support larger flocks (many urban roosts do not have enough trees for large flocks).
- » The roosts include food resources within the birds' daily foraging range (up to 6km from roosts).



Day roost habitat and crèche trees

As well as night roosts, black cockatoos need day roost habitat. Once daytime temperatures become too warm, black cockatoos must stop their foraging activity and take shelter in trees with shady canopies, where they rest until temperatures fall. Flocks need enough trees with dense canopies near their foraging areas, to allow them to rest and avoid heat stress on hot days. Likewise, black cockatoos benefit from access to a few trees with dense canopies near their roosts, to act as 'nursery' or 'crèche' trees when the flocks have new young with them. For example, a researcher has observed that when Carnaby's cockatoos return from their breeding sites to the Perth-Peel region in summer and early autumn each year, bringing their dependent young with them, they may crèche their young in one or more nursery trees near their roost each morning, while the adults head out to forage. The young birds retreat into the canopy to hide and rest, and wait to be collected and fed by their parents at the end of the day. They then return together to the night roost.

Actions »

Retain existing night roosts

1. **Identify** and protect all night roosts within the City of Kwinana. DBCA and BirdLife Australia have roost location data; BirdLife Australia will have the latest data if DBCA databases have not been updated.
2. **Work** with the WALGA and state government to enable local governments to protect important trees on private property.
3. **Recognise** that a roost may not be in use year-round but may be critical at particular times of year, related to seasonal availability of food nearby. Given the scarcity of food for black cockatoos in some areas and at some times of year, whenever a patch of foraging habitat is producing food, the roosts nearby will become critical. Without them, black cockatoos cannot access that food.

Create new roost habitat

1. **Plan** for roost habitat succession by planting new roost habitat.
2. **Recommended** roost trees in the Perth-Peel region include marri, jarrah and tuart. Over time, these will provide both roost habitat and food or corridor habitat. For Carnaby's cockatoos in the Perth-Peel region, pine also provides important roost habitat and food.
3. **Plant** and maintain several trees together where possible, to create a roost with enough trees to support larger flocks. Parks and schools are often appropriate locations to plant larger trees.
4. **Ensure** new roost trees are supported by enough food resources within 6km; preferably closer.
5. **Consider** planting new roost trees at or near an existing roost, to enable that roost to remain viable over several generations of trees.

6. **Ensure** that roost succession planning is supported by appropriate City strategies, policies and plans.
7. **Check** roost datasets (DBCA, BirdLife Australia) every 1–2 years, to ensure any new roosts are identified and protected.
8. **Notify** BirdLife Australia if new roosts are reported to the City of Kwinana, so they can update their database (greatcockycount@birdlife.org.au).

Community engagement

1. **Liaise** with BirdLife Australia to organise flock counts at roosts in the City of Kwinana through the annual citizen-science Great Cocky Count. Host and promote workshops about the Great Cocky Count, and the ongoing Cockywatch surveys (BirdLife Australia), to enable City of Kwinana community volunteers to become trained as citizen scientists. Direct funds towards Friends groups or other community groups to create new foraging habitat near roosts or enhance existing habitat (through infill planting; weeding).
2. **Provide** information through the City's online networks about the importance of retaining the City of Kwinana's black cockatoo roosts, and of providing enough food resources near roosts to ensure they remain viable.

Key points

- » **Protect all remaining night roosts for black cockatoos in the City of Kwinana.**
- » **Support roosts by providing food resources and clean drinking water nearby.**
- » **Create new roosts (succession planning) by planting marri, jarrah and tuart around ovals or in other public open spaces.**



The value of street trees

Along with their aesthetic value and urban cooling effects, mature street trees are valuable wildlife corridors. Murdoch University's flock tracking data show that one Carnaby's cockatoo flock in Perth's northern suburbs was able to access a patch of fruiting banksia on the far side of a dense new housing development by moving along a row of large tuart trees in the backyards of adjacent private properties, which had been protected by local government. No other vegetation had been retained in the vicinity, on either government or private land. Without the row of street trees, the black cockatoos may not have been able to access the important food resource on the other side of the development. In urban areas where food resources are already scarce, vegetation corridors are critical to allow black cockatoos to access all remaining foraging habitat.

1.3 Lack of vegetation corridors

- » In the Perth-Peel region, urbanisation has removed most black cockatoo habitat, and the habitat that remains has been fragmented into many smaller, unconnected patches.
- » Perth's urban landscape includes large expanses without trees, including many housing estates, industrial areas and some suburban blocks. Areas without trees are a threat to birds and other urban wildlife, because they are 'roadblocks' which prevent wildlife movement between natural areas, such as between one foraging area and the next.
- » Contrary to popular belief, when black cockatoos are resident in the Perth-Peel region, they do not just 'fly wherever they want'. Tracking data by Murdoch

University show that the birds avoid larger cleared urban areas without trees. Instead, they move through the landscape by relying on vegetation corridors, such as a row of street trees or a network of neighbouring habitat patches.

- » Patches and strips of vegetation offer black cockatoos protection, shelter, and places to land, forage and rest. They act as 'stepping stones', allowing the birds to move through built-up areas by making short 'step' flights between vegetation patches.
- » Without a vegetation corridor, food resources may not be accessible to black cockatoos.
- » Risks to vegetation corridors include the possibility that many trees and vegetation patches which form part of a corridor may be on private land, and not protected from clearing.





Actions »

- 1. Retain** vegetation corridors, to allow black cockatoos to access roosts and food resources.
- 2. Recognise** that patches of vegetation are valuable as part of a corridor even when each patch is small; including trees around ovals and parks and on private property. These are important to retain.
- 3. Undertake** succession planning for vegetation corridors. Where existing corridors comprise older trees, plan for retention of this resource for black cockatoos by planting new trees sufficiently far in advance to allow them to replace the older trees over time, enabling the corridor to remain. This will require long-term planning, and may require amendments to relevant City strategies, policies and plans.
- 4. Enhance** existing vegetation corridors and create new corridors, particularly where larger gaps exist between patches of foraging habitat.
- 5. Plant** patches of black cockatoo food trees (e.g. marri, jarrah, banksia, hakea) in spaces where there is little or no vegetation, and within 6km of roosts and food resources. New patches which can act as ‘stepping stones’ between larger areas of foraging habitat are particularly important.
- 6. Ensure adequate set-back distances when planting black cockatoo food trees on road verges. In quiet streets (speed limits <50km) plant at least 3–5m from road edges; on busier roads (speed limits >50km/hr) plant >10m from the road. Never plant black cockatoo food plants on median strips.**
- 7. Engage** the community to create vegetation corridors for black cockatoos in the City of Kwinana that will also provide food for the birds, including by planting food trees around school ovals, golf courses and in backyards. Assist Friends of Bushland groups to retain and enhance vegetation corridors, and support and promote community involvement in native vegetation planting days.

Key points

- » Black cockatoos need vegetation corridors to allow them to move through urban landscapes.
- » Planting new food resources can also increase vegetation corridors across the City of Kwinana.

Policy implementation

Creating and enhancing vegetation corridors for urban wildlife in the City of Kwinana is an effective way to implement the state Native Vegetation Policy, which includes a Strategic Goal of increasing the connectivity of native vegetation. It will also support local biodiversity and climate mitigation.



1.4 Lack of breeding habitat

Black cockatoos nest in large hollows that form in old eucalypts. To develop suitable hollows, the trees typically need to be at least 200 years old. Breeding habitat also needs to be supported by access to reliable fresh drinking water (preferably within 1km) and sufficient food (within 12km or preferably 6km). Black cockatoos prefer to breed near other members of their flock, so numerous hollows are required to support all breeding pairs.

Very little black cockatoo breeding habitat remains in the Perth-Peel region. In areas where breeding occurs but which have lost most of their natural hollows, artificial hollows have shown some success to date (largely for Carnaby's cockatoos), but artificial hollows require monitoring and maintenance over many decades to remain viable and safe. Hollows in poor repair have been linked to bird deaths (Carnaby's Cockatoo Recovery Team pers. comm). In urban areas, artificial hollows are often taken over by smaller parrot species such as corellas and galahs, which can become pests. Urban environments are also high-risk for black cockatoo fledglings, which are inexperienced fliers and vulnerable to vehicle strike and raven attack.

For the above reasons, the Department of Biodiversity, Conservation and Attractions (DBCA) advises against installation of artificial hollows in urbanised areas of the metropolitan Perth-Peel region.⁵⁶ This Plan supports that advice. In order to breed, the black cockatoos that roost and forage in the City of Kwinana will travel to breeding sites elsewhere at the start of the breeding season, and will bring their fledglings back to the City of Kwinana at the end of the breeding season.



Figure 10. Locations of confirmed Carnaby's cockatoo breeding activity (12km buffers)

Breeding records are only publicly available for Carnaby's cockatoos. To check whether parts of the City fall within 12km of confirmed or unconfirmed (i.e. potential) breeding locations for other black cockatoo species, please contact DBCA or BirdLife. The 12km buffer indicates the area within the City where food resources are supporting the breeding birds (Carnaby's cockatoos require sufficient food within 12km or ideally 6km of their nest).

Source: BirdLife Australia. Figure produced using Google Earth Pro (<https://earth.google.com>).

Note: High resolution maps are available. See page 5 of this document for more information. 

Although no known natural breeding sites remain in the City of Kwinana, parts of the City are within the 12km foraging buffer areas around the locations of confirmed Carnaby's cockatoo breeding activity (Figure 10). Carnaby's cockatoos rely on foraging habitat within 12km of their nests, to feed themselves and their young during the breeding season.

All foraging habitat in the City of Kwinana that lies inside these 12km buffers should be considered an important food resource for Carnaby's cockatoo breeding pairs and their young.

⁵⁶ For further information, see DBCA's 'Fauna Note Artificial Hollows for All Black Cockatoos': www.dbca.wa.gov.au/management/threatened-species-and-communities/resources/threatened-and-priority-fauna-resources

Actions »

Do not install artificial hollows. Black cockatoo breeding is not recommended in urban areas of Perth and Peel.

Support breeding success

The City of Kwinana can support the breeding success of its black cockatoos by providing enough food resources during the non-breeding season to enable black cockatoo pairs to build body condition, which will help them breed successfully when they travel to their breeding sites outside the City. For Carnaby's cockatoos that are breeding nearby and relying on the City's food resources to help their young survive, the City can focus on protecting and enhancing foraging habitat within the buffer areas identified in **Figure 10**. See **Table 1** for food species to plant.

Support young black cockatoos

The City of Kwinana can support the survival of the young black cockatoos that arrive back in the City with their parents and flock at the end of the breeding season, by providing sufficient quantities and diversity of food plants for the parent birds while they are feeding their young, and for the young birds as they learn to forage. Roosting habitat, reliable fresh drinking water and vegetation corridors (see earlier sections of this Plan) will also help black cockatoos to breed and their young to survive.

Manage nest competitors

- » Smaller, more common parrot species such as corellas can build up in urban areas and then travel to breeding areas outside the city, where they can compete with black cockatoos for scarce nesting hollows. For guidance on managing common parrot species in urban areas so



that they do not become overabundant, contact BirdLife Australia or the state Department of Primary Industries and Regional Development (DPIRD). As of 2023–24, DPIRD is developing an official strategy for managing overabundant parrot species.

- » Feral European honey bees are a major threat to black cockatoo breeding success and conservation. When European bees are allowed to swarm away from their hives and become feral, they can take over tree hollows that were used by black cockatoos or other wildlife. In some areas, many hollows have been taken over by bees, making them unavailable for black cockatoos. Black cockatoo chicks have been found either stung to death or suffocated by feral bee swarms.⁵⁷ Local governments in urban areas can support black cockatoos by raising awareness of the need for owners of bee hives to ensure their bees do not swarm away from the hive and become feral.

Crèche trees for fledglings

For the first few weeks after returning from their breeding sites, when young black cockatoos are particularly vulnerable, some birds may 'crèche' their dependent young in a suitable tree near their roost, to shelter during the day while the flock forages. The flock then collects the young birds before returning to the roost to sleep. The City of Kwinana could support its black cockatoo flocks by identifying a quiet green space within 1–2km of each roost, in which to plant a few trees that will grow to become medium to large trees with relatively dense foliage, to act as crèche or nursery trees. Macadamia trees are ideal, as they provide both dense foliage for young birds to hide in, and high-energy food. If the City of Kwinana is installing some Cockitrough bird waterers (see **Lack of fresh drinking water** section of this Plan), planting crèche trees at the same location as the bird waterers could be ideal, as it allows flocks to drink and crèche their young at the same time, freeing up more of the day for foraging.

PHOTO Rick Dawson



⁵⁷ Research conducted by the WA Museum: <https://museum.wa.gov.au/explore/online-exhibitions/cockatoo-care/feral-bees>

1.5 Tree diseases

- » Black cockatoo food resources and roosting habitat are at risk from a number of plant diseases. The national Carnaby's Cockatoo Recovery Plan identifies a range of pathogens that can kill black cockatoo habitat, including leaf and canker pathogens (e.g. marri canker), and soil borne pathogens, such as the highly aggressive root rot pathogen *Phytophthora cinnamomi*.

- » Phytophthora root rot disease or 'dieback' is a particular threat to black cockatoos. The pathogen is difficult to remove once it is in the soil and can cause devastating losses in banksia and other Proteaceous habitat, as well as jarrah – all critical resources for Carnaby's cockatoo survival.
- » In the Perth-Peel region, particular vigilance is required to manage the health of black cockatoo food resources, because most native habitat in the region has been reduced to small, fragmented patches. Small patches are more vulnerable than larger patches to disease being introduced from outside.



1.6 Habitat loss from fire

- » Fire in native vegetation can cause large direct losses of black cockatoo food resources, and also (depending on the site) roosts and breeding habitat.
- » Black cockatoo habitat in the Perth-Peel region is at particular risk from deliberately-lit fires, given that most habitat in the region is readily accessible.
- » The risk to black cockatoo habitat from fire is exacerbated by climate changes. The Perth-Peel region is now experiencing hotter temperatures, less rainfall, and other harmful climate changes which are making fires hotter and more destructive to black cockatoo habitat.

Actions »

- 1. Prioritise** monitoring and management of the health of the City of Kwinana's native vegetation and follow best practice in disease prevention.
- 2. Ensure** the City and its contractors have appropriate hygiene policies and practices to prevent dieback disease from entering new areas of native vegetation in the City. Common ways that the pathogen can be introduced include via contaminated soil, mulch and fill products (including during planting), and on shoes, tools, tyres and other vehicle components. Use gardening materials accredited with the Nursery Industry Accreditation Scheme Australia (NIASA accredited products)⁵⁸.
- 3. Ensure** the City and its contractors have appropriate hygiene policies and practices to manage marri canker e.g. sterilising pruning tools.

- 4. Plant** marri where possible, recognising the City's critically endangered Baudin's cockatoos depend on marri. Although marri is at risk from canker, it is important to plant marri and manage the disease risk effectively; without marri, the City's Baudin's cockatoos will be lost.
- 5. Ensure** the City's strategies, policies and plans, wherever relevant, are developed or updated to enable effective health monitoring and disease prevention in the City of Kwinana's native vegetation.
- 6. Consult** state and Commonwealth guidelines on management of dieback and other native plant diseases as required. BirdLife Australia and natural resource management organisations operating in the Perth-Peel region also provide advice.

Actions »

- 1. Monitor** and manage the risks from fire using evidence-based approaches; such as raking⁵⁹.
- 2. Manage** the increasing risks of high-intensity fires in the Perth-Peel region by ensuring the City of Kwinana becomes a local government leader in climate change mitigation (see [Climate change and Urban Heat Island effects](#) section).

⁵⁸ BirdLife Australia has a printable pamphlet on the topics of choosing the best food plants for black cockatoos and managing the risk of dieback in native trees: www.calameo.com/read/0041078955a20c96b51c8

⁵⁹ Raking of leaf litter from around the bases of important trees (e.g. large trees with hollows; roosts) helps them survive fire; including prescribed burns. Contact DBCA for specific distance to rake from the base and other advice.

Threat 2. Lack of fresh drinking water

Lack of access to fresh drinking water is a threat to black cockatoos across the Perth-Peel region. Risks and impacts include heat stress, disease from contaminated water sources, and vehicle strike when drinking from pooled water on roads and verges.

2.1 Heat stress

- » Without water, black cockatoos can die quickly from heat stress on very hot days. Losses of almost entire flocks have occurred due to heat stress. This risk will increase as Perth-Peel becomes hotter, with the region now experiencing record-breaking heat.
- » Black cockatoos have specific water requirements.
 - They need access to fresh water twice daily, in the morning and late afternoon, as they leave and return to their roosts.
 - They need the fresh water source to be as close as possible to their roost, preferably within 1km, and in a safe location.



Heat stress avoidance strategies

As well as drinking water every morning and evening, black cockatoos avoid dangerous overheating by refraining from flying and foraging once temperatures become too warm, and retreating into trees with heavy shade. Carnaby's cockatoos remain inactive for as long as temperatures remain over 30°C (Saunders 1977). New Murdoch University research shows that for every 5°C increase in maximum temperature, Carnaby's cockatoos need to shelter in place for an additional 30 minutes.⁶⁰ This need to shelter during hot weather reduces the time available to find enough food; highlighting the risks from climate change.

- » Botulism outbreaks can occur in ponds and lakes in Perth and Peel. The bacterium that produces botulism toxin is often present in standing water; but is only triggered to produce the toxin in certain environmental conditions, when oxygen levels in water are low (e.g. in warm weather with low water levels or build-up of organic matter). Botulism outbreaks can occur repeatedly if water quality is not managed. The disease is often fatal in birds, and can kill large numbers. Waterbirds are most commonly affected, but any bird drinking the water is at risk.

2.3 Vehicle strike when drinking on roads and verges

- » As black cockatoos prefer to drink from shallow rather than deep water, in the absence of reliable drinking water sources they are attracted to drink from rainwater or sprinkler runoff that has pooled in potholes and on roadsides without adequate drainage. This leads to high risk of vehicle strike. See [Vehicle strike](#) section.



2.2 Poor water quality

- » Black cockatoos have a natural preference to drink from shallow water, rather than deeper water bodies such as lakes, and this can expose them to poor quality water. They will drink from pond-edges, troughs and bird baths, water that has pooled in potholes and beside roads without adequate drainage, and even water that collects in leaves of leafy vegetables in market gardens.
- » In the Perth-Peel region, poor-quality and contaminated water sources have been linked to disease outbreaks and deaths of black cockatoos.
- » In 2023, an outbreak of black cockatoo illness and deaths occurred in Perth's northern suburbs. Investigations by Murdoch University and Perth Zoo Veterinary Department are ongoing, however chemicals were detected in the gut contents of deceased birds, including pesticides; suspected to have been ingested from contaminated drinking water.

⁶⁰ Riley et al. (2023).

Actions »

Establish a network of bird water drinking stations

This will provide all black cockatoo flocks in the City of Kwinana with safe, reliable access to fresh drinking water close to their roosts.

What to install

- » Artificial waterers for black cockatoos have specific design requirements for effectiveness and safety. A leading design is the Cockitrough, designed and supplied by the Town of Victoria Park.
- » The Cockitrough features several shallow-water drinking platforms to allow several birds to drink at once, which aligns with natural black cockatoo behaviour. Troughs are self-flushing and situated high off the ground to protect birds from cats and dogs. In 2024, the City of Kwinana collaborated with the *Keep Carnaby's Flying – Ngoolarks Forever* project, to purchase two Cockitroughs and install them near known roosts.
- » Information on the Cockitrough is available from the Town of Victoria Park: www.victoriapark.wa.gov.au/Around-town/Environment/Bird-waterers

Where to install:

- » Prioritise providing water at key (i.e. large) roosts (nearby lakes and ponds are not appropriate drinking water sources, given the risk of poor water quality and given that black cockatoos prefer to drink from shallow water). Roost location and flock size data are available from DBCA and BirdLife Australia; BirdLife is also available to discuss location options and provide advice (Carnabys@birdlife.org.au).



- » Install the water drinking station at the roost site or at the closest suitable location, preferably within 1km.
- » To avoid health risks for the birds, water provided at drinking stations should be potable water. Cockitroughs can be easily installed at sites with scheme water, or as an off-grid system using an intermediate bulk container (IBC) which can be filled periodically.
- » Install close to shaded trees or other shaded perch options, to protect birds from heat stress on very hot days while they take turns drinking, and to protect the 'sentinel birds' which act as lookouts for the flock.
- » Install in a quiet location away from traffic, carparks and too much human or dog activity.

Cockitrough' bird waterers installed in Jirdarup Bushland, Victoria Park, being used by a flock of forest red-tailed black cockatoos. Note the importance of installing waterers near trees, which offer shade and perches for flock members as they wait their turn.

PHOTO Georgina Wilson

- » Cockitrough suppliers will have additional information about location requirements. BirdLife Australia offers expert advice about appropriate locations (Carnabys@birdlife.org.au).
- » Establish a network of drinking stations across the City, so all roosts have a source of safe fresh drinking water.
- » Engage the community and local businesses to raise funds for Cockitroughs.

Maintenance of Cockitroughs

- » Cockitroughs are designed to be low maintenance, with self-flushing to replenish clean water at chosen intervals (in warm weather, adjust the setting to replenish water more often). Important maintenance includes checking that the trough is filling correctly (check at least every 2 months; more frequently in hot weather), checking batteries (depending on valve type), and annual visual inspection (in Sept-Oct) to remove any heavy debris.

Manage water quality in ponds and lakes

- » Monitor and manage water quality in ponds, lakes and wetlands, including risks of chemical contaminants and disease outbreaks from bacteria and algal blooms.
- » Manage botulism outbreaks.
- » Maintain records of water quality in individual ponds, lakes and wetlands to help with water resource management.
- » Where a pond, lake or wetland has repeated water quality issues and is within 1km of a black cockatoo roost, prioritise installation of a Cockitrough at that roost to provide a safer option.

Note: as noted earlier, the presence of ponds and lakes does not mean these are appropriate sources of drinking water for black cockatoos. There may be accessibility issues, and water quality may be poor. It is preferable to provide a network of drinking stations.

As part of this Conservation Action Plan, roost data and locations of green space in the City of Kwinana were analysed to identify priority locations for installation of Cockitrough water drinking stations (Figure 11). These locations take into account that some important roosts may already have a bird waterer installed nearby.




Figure 11. Recommended locations to install Cockitrough water stations

Priority locations have been identified based on proximity to known key night roosts and overall site suitability, to create a network of safe watering points across the City. Where possible, locations are at the roost or within 1km. Where no suitable City-managed green space is available, sites have been identified in the closest Bush Forever area.

From north to south, recommended sites are: Bruny Meander Park, Wandí; Orton Road Bush Forever site (Casuarina Prison), Casuarina; Glenworth Drive, Wellard; and Henley Park, Wellard.

Note: while these locations are considered ideal due to roost proximity and overall site suitability, if installation is not feasible logistically at a site, an alternative location should be sought, close to the recommended location.

Source: BirdLife Australia and Murdoch University. Green space data as per Figure 5. Figure produced using Google Earth Pro (<https://earth.google.com>).

Reduce access to drinking water on and beside roads

- » Maintain roads and verges to prevent potholes and improve water drainage away from roadsides. Maintain water infrastructure near roads to avoid water leaks. Redirect sprinklers away from roads. For further details, see actions listed under **Vehicle strike**.

These actions to address the threat to black cockatoos from lack of fresh drinking water will be relevant to feed into many of the City's strategies, policies and plans, such as those related to urban biodiversity, climate adaptation, and roads.

Key points

- » Black cockatoos need drinking water daily, close to their roosts.
- » They will drink from potholes and roadsides after rain. Repair potholes and road drainage to prevent vehicle strike.
- » Poor water quality has resulted in illness and death of black cockatoos in the Perth-Peel region.
- » Cockitroughs or similar designs provide safe potable water (see previous page).

Note: High resolution maps are available. See page 5 of this document for more information.



Threat 3. Vehicle strike

After loss of habitat, vehicle strike is one of the most significant threats to black cockatoos throughout their distribution range, particularly in high-traffic areas and on roads where vehicles are travelling at speed⁶¹. Vehicle strike causes traumatic injuries ranging from minor wounds and bruises to catastrophic bone fractures.

Common vehicle strike injuries include keel fractures, wing fractures, concussions, and trauma causing internal bleeding. Many black cockatoos that are hit by cars will die without being found. Even among those retrieved and admitted for treatment, around 40% will have injuries so severe that they are euthanased. Black cockatoos are at risk of vehicle strike when:

- » feeding on low to medium-height vegetation on roadsides and median strips, as the birds will be at vehicle height when flying in and out of these food resources.
- » feeding on grain spill (on roads used by grain haul vehicles).
- » drinking from potholes in roads or puddles that have formed in roadside depressions.

Vehicle strikes in the City of Kwinana

To support the development of the City of Kwinana Black Cockatoo Conservation Action Plan, Murdoch University undertook an epidemiological analysis to identify the causes of admission to the Perth Zoo Veterinary Department (PZVD) for black cockatoos from the City of Kwinana 2018–23. Vehicle strike was one of the two main identified causes of admission (**Figure 12**. See also **Appendix 2**).




Figure 12. City of Kwinana black cockatoo admissions to Perth Zoo Veterinary Department 2018–23: Reported vehicle strikes

Between 2018–23, 10 black cockatoos from the City of Kwinana were admitted to PZVD with evident road strike injuries: five Carnaby’s cockatoos and five forest red-tailed black cockatoos. Several other cases were admitted with ‘unknown cause’; many of these are also likely to be vehicle strike cases that were momentarily stunned or had minor concussions (**Appendix 2**). These cases represent a minimum, as many birds with vehicle strike injuries may not be found or reported.

Most reported vehicle strikes were in the first half of the year (**Figure 12**), likely due to the higher numbers of Carnaby’s cockatoos in the City of Kwinana at this time of year, when migrating flocks return from breeding sites to spend the non-breeding season in this area.

61 Le Souëf, A. et al. (2015). Presentation and prognostic indicators for free-living black cockatoos (*Calyptorhynchus* spp.) admitted to an Australian zoo veterinary hospital over 10 years. *Journal of Wildlife Diseases*, 51(2), 380–388.

Actions »

- 1. Apply** set-backs from roads: To ensure black cockatoos have enough food in urban areas, it is important to plant food trees, rather than non-food trees, whenever possible. Road verges often present an excellent opportunity. When planting food trees near roads, apply set-backs to avoid vehicle strike:
 - in quiet suburban streets (speed limit <50km/hr), plant food trees at least 3–5m back from road edges. Plant food trees on a single side to reduce the risk of birds flying across the road between trees⁶².
 - for busy roads (speed limit higher than 50km/hr), plant food trees at least 10m from road edges.
 - maintain existing large food trees (jarrah and marri) on verges by pruning low branches.
 - never plant black cockatoo food plants on median strips; this is a death trap for the birds.
- 2. Plant** and maintain areas of high-quality food resources in areas away from roads, to attract flocks to those areas.
- 3. Maintain** road surfaces and verges to avoid pooling water which attracts black cockatoos:
 - monitor and fill potholes and depressions.
 - ensure road water runoff drains well away from verges.
- 4. Maintain** council-managed water infrastructure near roads (taps, pipes, hoses) to prevent water leaks.
- 5. Direct** verge sprinklers away from roads.



- 6. Maintain** records and build a dataset of vehicle strike locations. If hotspots for vehicle strike are identified, particularly repeated strikes at the same location, investigate and undertake mitigation actions if required, and consider installation of road signage.
- 7. Report** injured black cockatoos to the volunteer-run Black Cockatoo Rescue Service at **Kaarakin Black Cockatoo Conservation Centre: (08) 9390 2288** during work hours or **0448 046 202** after hours, or Parks and Wildlife (DBCA) Wildcare Helpline: **(08) 9474 9055**.

- 8. Support** black cockatoos with community engagement in the City of Kwinana, highlighting 'What to do if you find an injured black cockatoo' on the City's website, including providing links to the Black Cockatoo Rescue Service at **Kaarakin Black Cockatoo Conservation Centre** and the **Found an injured bird?** page on the *Keep Carnaby's Flying – Ngoolarks Forever* website.

⁶² Peck A. & Stokes V. (2019). *Black-cockatoo conservation and recovery: Guidelines for councils and land managers*. BirdLife Australia, November 2019.

Figure 13 shows the reported locations of vehicle strike cases in the City of Kwinana, for black cockatoos admitted to Perth Zoo Veterinary Department 2018–23. This map is produced to support the City’s record-keeping and monitoring of this threat over time, and to enable investigation of any issues near that location that could predispose black cockatoos to vehicle strike (e.g. potholes, road drainage issues, or food resources next to the road).

Key points

- › Fix potholes and roadside drainage to deter birds from drinking on and near roads.
- › Apply setbacks from roads when planting food trees.




Figure 13. Reported locations of vehicle strike cases in the City of Kwinana, for black cockatoos admitted to Perth Zoo Veterinary Department 2018–23

Note: High resolution maps are available. See page 5 of this document for more information.



BLACK COCKATOO CONSERVATION



CHiPS-affected Carnaby's cockatoos can often be rehabilitated and returned to the wild if they receive veterinary care.

PHOTO Rick Dawson

Threat 4. Carnaby's Hindlimb Paralysis Syndrome

Carnaby's Hindlimb Paralysis Syndrome (CHiPS) has been recognised in Carnaby's cockatoos since 2012⁶³. To date Carnaby's cockatoos are the only species known to be affected.

Carnaby's cockatoos with this syndrome are admitted to Perth Zoo Veterinary Department (PZVD) in the summer and autumn months with a weakness or paralysis in their hindlimbs, thought to be a delayed-onset neurological disease caused by pesticide exposure. Some black cockatoos with CHiPS are too sick to be treated and require euthanasia, but most can be treated with supportive care and released back to the wild. CHiPS-affected Carnaby's

cockatoos tend to present to PZVD having been found in urban areas where there are key roost sites that support relatively large numbers of birds, and from areas where there are more people to notice the affected birds.

Research into this syndrome is ongoing, however it is thought the birds may be exposed to pesticides in agricultural areas outside Perth during the breeding season and, given the suspected delayed onset of this condition, are noticed in a debilitated condition when they return to their urban roosts and foraging areas at the start of the non-breeding season.

For the period of investigation there were no CHiPS cases admitted to PZVD from within the City (**Appendix 2**). However, it is important to note that CHiPS cases may go unnoticed and unreported; particularly in peri-urban and rural areas with fewer people to notice and report debilitated birds.

Actions »

- 1. Mitigation** measures for CHiPS are difficult to identify at present and require further research into the source(s) of the toxin. However, awareness of the condition is important, as although the affected birds will die if left in the wild, many can be successfully rehabilitated if taken into care. Sick or injured black cockatoos should be reported to the volunteer-run Black Cockatoo Rescue Service at **Kaarakin Black Cockatoo Conservation Centre: (08) 9390 2288** during work hours or **0448 046 202** after hours, or Parks and Wildlife (DBCA) Wildcare Helpline: **(08) 9474 9055**.
- 2. Support** black cockatoos with community engagement, highlighting 'What to do if you find an injured black cockatoo' on the City's website (see **Vehicle strike** section of this Plan for details).



63 Le Souëf, A., Vitali, S., Dawson, R., Vaughan-Higgins, R., & Warren, K. (2020). Hindlimb paralysis syndrome in wild Carnaby's cockatoos (*Calyptorhynchus Latirostris*): A new threat for an endangered species. *Journal of Wildlife Diseases*, 56(3), 609–619.

Threat 5. Raven attack

Raven attack is a threat to black cockatoos in urban areas which has emerged in the last decade and mainly affects forest red-tailed black cockatoos. The increase in raven attacks on forest red-tailed black cockatoos in urban areas has coincided with an increase in movement of flocks of this black cockatoo species into urban areas, starting in the mid-2000s, to forage on introduced Cape Lilac, in line with declines in their native food trees (marri and jarrah) due to habitat clearing, fire and impacts of climate change⁶⁴. There has also been an increase in ravens, a successful urban wildlife species, in Perth over the past 30 years.

Injuries caused by ravens tend to be around the tail end of the cockatoo, with a classic presentation being loss of tail feathers. There are sometimes wounds to the skin as well, and in some cases these wounds are too severe for treatment and require euthanasia. However, most raven-attacked black cockatoos can be treated and returned to the wild. 'Secondary' raven attacks sometimes occur when black cockatoos of any species are debilitated for another reason, such as vehicle strike or Carnaby's hindlimb paralysis syndrome. Secondary attacks further reduce these black cockatoos' chances of survival. Ravens are more numerous in areas populated by more people, because they benefit from the changes associated with urbanisation.



Manage food waste to deter ravens and protect forest red-tailed black cockatoos.

In the epidemiological analysis undertaken for this Conservation Action Plan by Murdoch University to examine the causes of admission to the Perth Zoo Veterinary Department for black cockatoos from the City of Kwinana 2018–23, raven attack was not one of the main reasons for admission (see **Appendix 2**).

Actions »

- 1. Manage** food waste to avoid attracting ravens.
 - Self-closing bin lids
 - Community awareness-raising regarding food waste management and the importance of not feeding ravens and other wildlife, including fact sheets; monitoring and compliance as required
 - If hotspots for raven attack are identified, investigate and reduce access to artificial food sources
 - As noted in the 'Living with ravens' Information Sheets produced by several councils, appropriate food waste management is the most effective measure for deterring ravens. Although raven culls have occurred (with relevant lawful authority) in the past in Perth-Peel to control raven numbers in problem areas, it is not clear if this is an effective measure to safeguard black cockatoos, and appropriate food waste management is recommended.
- 2. Report** sick or injured black cockatoos to **Kaarakin Black Cockatoo Conservation Centre: (08) 9390 2288** during work hours or **0448 046 202** after hours, or DBCA's Wildcare Helpline: **(08) 9474 9055**.
- 3. Support** black cockatoos with community engagement, highlighting 'What to do if you find an injured black cockatoo' on the City's website (see details in the **Vehicle strike** section of this Plan), and advising the public to report cases of black cockatoos that are being attacked by ravens on the ground and appear to be struggling.

⁶⁴ Johnstone, R.E., Kirkby, T., & Sarti, K. (2017). The distribution, status, movements and diet of the forest red-tailed black cockatoo in the south-west with emphasis on the Greater Perth Region, Western Australia. *Western Australian Naturalist*, 30, 193–219.

Threat 6. Illegal shooting

Black cockatoos are sometimes illegally shot, particularly when entering fruit or nut orchards in regional areas. Black cockatoos that are found with shooting injuries have not necessarily been shot locally, but may have travelled from other areas after being injured. As of 2023, fines for shooting black cockatoos are up to \$400,000 for individuals⁶⁵. Shooting injuries can include soft tissue trauma, as well as bone fractures and internal bleeding.

In the epidemiological analysis undertaken for this Conservation Action Plan by Murdoch University to examine the causes of admission to the Perth Zoo Veterinary Department for black cockatoos from the City of Kwinana 2018–23, records show that two shooting cases were admitted (**Figure 14** and **Appendix 2**).



⁶⁵ As per section 150(1) of the BC Act in relation to taking of threatened fauna [as of 2023].




Figure 14. Reported locations of shooting cases in the City of Kwinana, for black cockatoos admitted to Perth Zoo Veterinary Department 2018–23

These are sites where injured birds were recovered and are not necessarily where the shooting occurred.

Actions »

1. While there are no specific actions required to manage this threat in the City of Kwinana, it may be useful to **recognise** that black cockatoos which have been shot and injured elsewhere may occasionally be found debilitated in the City of Kwinana. Sick and injured birds should be reported to **Kaarakin Black Cockatoo Conservation Centre: (08) 9390 2288** during work hours or **0448 046 202** after hours, or DBCA Wildcare Helpline: **(08) 9474 9055**.

Note: High resolution maps are available. See page 5 of this document for more information.

Threat 7. Climate change and Urban Heat Island effects

Perth-Peel is experiencing higher temperatures, declining rainfall and other climate changes driven by use of fossil fuels and greenhouse gas emissions.

Climate threats are increasing in the absence of effective mitigation nationally and globally. Effective mitigation by government will require direct action at the local level, as well as community engagement, planning, advocacy and longer-term action by all levels of government.

The WALGA Advocacy Position Manual (2023) states the following:

Local Government acknowledges:

1. The science is clear: climate change is occurring and greenhouse gas emissions from human activities are the dominant cause.
2. Climate change threatens human societies and the Earth's ecosystems.
3. Urgent action is required to reduce emissions, and to adapt to the impacts from climate change that are now unavoidable.

WA Local Government is committed to addressing climate change [and calls for] strong climate change action, leadership and coordination at all levels of government.



Climate threats to black cockatoos in the Perth-Peel region include increasing risk of heat stress-related deaths (see [Lack of fresh drinking water](#) section) and loss of food resources from drought and higher-intensity fires (see [Habitat loss from fire](#) section).

Hotter temperatures and lower rainfall lead to poorer tree health and lower seed productivity for key black cockatoo food species, reducing food availability.⁶⁶

Additional climate-related risks include longer enforced non-feeding periods during the warmest parts of the day (once temperatures reach around 30°C), during which time black cockatoos will generally stop foraging and shelter-in-place in shady vegetation to avoid heat stress.⁶⁷ This shorter feeding time reduces their capacity to build energy reserves in the lead-up to the breeding season, or to feed nestlings during the breeding season.

⁶⁶ Government of Western Australia (2019) EPA Advice: Carnaby's Cockatoo in Environmental Impact Assessment in the Perth and Peel Region; Saunders et al. (2021) Carnaby's Black-Cockatoo *Zanda latirostris*. In *The Action Plan for Australian Birds 2020*.

⁶⁷ Saunders D.A. (1977) The effect of agricultural clearing on the breeding success of the White-tailed Black Cockatoo. These findings are supported by new Murdoch University flock tracking research: Riley et al. (2023). Accelerometry reveals limits to use of an energy-saving anthropogenic food source by a threatened species; a case of Carnaby's cockatoos (*Zanda latirostris*) and canola. *Ecology and Evolution*, 2023;13:e10598.



Plants that provide urban cooling and black cockatoo food together — an opportunity for local government:

- » Some local governments are planting trees for urban cooling and greening that are not appropriate for black cockatoos and other wildlife. This is a missed opportunity to support urban cooling, greening and black cockatoos together.
- » It is valuable to remember that the types of native trees and shrubs that are essential for black cockatoo survival, including banksia and hakea, support urban cooling and Urban Heat Island mitigation due to the plant's evapotranspiration, whereby water is transpired through the plant to the leaf surface and absorbs heat energy from the air as it evaporates, thereby cooling the air. (Taller food trees, i.e. marri, jarrah and non-local macadamia and pine, also provide shading; but all black cockatoo food plants, including banksia and hakea, assist urban cooling via evapotranspiration).
- » Lack of awareness of the role of evapotranspiration in urban cooling can lead people to believe that they need to plant tall shady trees to mitigate Urban Heat Island effects, when they could also plant black cockatoo food plants which cool the air through evapotranspiration. This supports urban cooling and greening while helping ensure the City does not lose its black cockatoos.
- » **Planting trees for urban cooling that specifically include black cockatoo food trees and shrubs will enable local government to achieve outcomes for urban cooling, urban forests, biodiversity and black cockatoo conservation in a single action.**


Actions »

1. **Undertake** actions to reduce the City of Kwinana's greenhouse gas emissions to net zero, to help avoid the worst impacts of urban heating for the people of Perth and the City's black cockatoos.
2. **Engage** and support City of Kwinana residents to reduce their carbon footprint, including promoting and supporting clean energy modes of travel and energy-saving/ clean energy options and supporting community tree-planting initiatives.
3. **Prioritise** planting of black cockatoo food species wherever possible, when planting for urban cooling. Recognise that planting native trees and shrubs will mitigate Urban Heat Island effects and promote urban cooling even without shading effects; the urban cooling will occur due to the native vegetation's evapotranspiration capacity. This includes evapotranspiration from black cockatoo food trees and shrubs such as banksia and hakea (even small patches), as well as larger trees (marri, jarrah).
 - Adopting a strategy of “net gain” of black cockatoo food resources will meet urban forest / canopy goals and reduce Urban Heat Island effects while simultaneously supporting black cockatoos and local biodiversity.
4. **Encourage** City of Kwinana residents to plant gardens (preferably with black cockatoo food species). Educate the community that trees, bushes, shrubs and even vegetable garden beds in private gardens have much higher evapotranspiration and urban cooling capacity than lawns.⁶⁸ Compared with lawns, native gardens that include black cockatoo food plants (see **Table 1**) also provide food for other wildlife and help to conserve Perth's increasingly limited water.

⁶⁸ Small, G., Jimenez, I., Salzl, M., & Shrestha, P. (2020). Urban heat island mitigation due to enhanced evapotranspiration in an urban garden in Saint Paul, Minnesota, USA. *WIT Trans. Ecol. Environ.*, 243, 39–45.

5. **Engage** with and support local 'Friends of bushland' community groups, to optimise the ability of these groups to provide additional capacity, alongside the City's staff, to plant native vegetation that will support both black cockatoos and urban cooling.
6. **Look** for opportunities to simultaneously take action for urban cooling and for the City's black cockatoos. For example, the WALGA provides information and resources about climate change for local governments. When accessing and using this information, look for possibilities to simultaneously take action for black cockatoos, such as by leveraging the urban cooling capacity of black cockatoo food trees.

7. **Advocate** to the WALGA for Urban Greening to explicitly support the planting of black cockatoo food trees, including banksia, hakea, marri and jarrah, in recognition of the capacity of this native vegetation to mitigate Urban Heat Island effects through evapotranspiration (as well as shading, for marri and jarrah). Promotion of Urban Greening actions that address both urban cooling and black cockatoo conservation (local biodiversity) will be critical in urban areas, where limited space exists for tree planting, and where addressing several outcomes simultaneously will support local governments to meet key climate and biodiversity goals.
8. **Update** the City's strategies, policies, plans and guidelines, where required, to support the above actions for climate change mitigation, urban cooling and concurrent black cockatoo conservation.



Key points

- » Net zero greenhouse gas emissions are critical to avoid the worst impacts of urban heating.
- » Planting black cockatoo food trees will contribute to urban cooling through the trees' evapotranspiration, while helping address food scarcity for black cockatoos.

Threat 8. Cumulative impacts

In the past, and in some cases still today, local government planners and decision-makers have neglected to consider cumulative impacts, when considering the impacts of decisions and actions on threatened black cockatoos.

It is this failure of planners and decision-makers to recognise the cumulative harm from many individual actions that has allowed the ongoing 'net habitat loss' in local government areas, thereby forcing black cockatoo populations to continue to decline.

There is currently a large number of development actions and associated clearing of remnant black cockatoo habitat across the Perth-Peel region. Many clearing actions involve clearing of small patches of important habitat or individual important trees that do not individually meet the threshold for requiring referral or controlled action (under the federal EPBC Act), but which together involve the loss of extensive areas of foraging, breeding and/or corridor habitat used by black cockatoo populations.

The risks and impacts posed to the viability of black cockatoos from cumulative habitat losses are, at present, greatly under-appreciated. The likely impacts become even greater when the impacts from other individual sources of harm, including vehicle strikes, disease and destructive climate changes, are added to the cumulative impact total.



A Carnaby's cockatoo flock flies over a cleared area that used to contain food resources and roosts.

Actions »

- 1. Ensure** all planning and decision-making related to black cockatoo habitat and conservation involve explicit consideration of the risks from cumulative impacts, including future risks, and management of these risks.
- 2. Ensure** the City's planning and regulatory decisions and actions do not lead to net loss (and preferably, achieve net gain) of food resources for black cockatoos in the City of Kwinana. Success regarding net gain of black cockatoo foraging habitat could be reported annually on the City's website.
- 3. Advocate** to state government, directly or through the WALGA, for additional local government powers to protect significant trees on private property, to help local government to manage the cumulative impacts of habitat loss on black cockatoos and other threatened species. Similarly, advocate for additional state government resources to support the evaluation and management of cumulative impacts on urban biodiversity, including the City of Kwinana's flocks of Carnaby's cockatoos and forest red-tailed black cockatoos.


Key points

- › Impacts to black cockatoos may appear insignificant when considered separately, when cumulatively they represent 'death by a thousand cuts'.
- › Proper consideration of cumulative impacts from all sources is essential to safeguard the City of Kwinana's black cockatoos for current and future generations.

Actions table: Summary of actions to safeguard black cockatoos in the City of Kwinana

This table summarises the key threats to black cockatoos in the City of Kwinana, and actions for the City to undertake to retain its flocks of Carnaby's cockatoos and forest red-tailed black cockatoos. All actions are suitable to develop into SMART-style goals. For details about each action, refer to the relevant section of the Plan, including maps to support the implementation of specific actions.

Threat	Actions	Page	
1 Loss of habitat	Lack of food resources For details of this threat, see page 12	Retain and improve existing food resources	page 16
		Create additional food resources	page 18
	Lack of roosts For details of this threat, see page 29	Retain existing night roosts	page 31
		Create new roost habitat	page 31
		Engage the community	page 31
	Lack of vegetation corridors For details of this threat, see page 32	Retain existing vegetation corridors	page 33
		Create new corridor habitat	page 33
		Ensure safe set-back planting distances from roads when planting black cockatoo food trees	page 33
		Engage the community	page 33
	Lack of breeding habitat For details of this threat, see page 34	Support breeding success	page 35
		Support young black cockatoos	page 35
		Manage nest hollow competitors (feral European honeybees; smaller parrots)	page 35
	Tree diseases For details of this threat, see page 36	Monitor and manage native vegetation health	page 36
		Ensure effective hygiene practices to prevent dieback	page 36
		Update the City's strategies, policies and plans as required	page 36
	Habitat loss from fire For details of this threat, see page 36	Manage direct risks	page 36
Manage indirect risks from climate changes/urban heating		page 36	



Summary of actions to safeguard black cockatoos in the City of Kwinana (cont.)

Threat			Actions	Page
2	Lack of fresh drinking water	For details of this threat, see page 37	Establish a network of water drinking stations	page 38
			Manage water quality in ponds and lakes	page 39
			Reduce access to drinking water on/beside roads	page 40
3	Vehicle strike	For details of this threat, see page 41	Apply planting set-backs from roads when planting food trees. For safe set-back distances depending on road speed limits	page 42
			Maintain road surfaces and verges to avoid pooling water (water drinking hazard)	page 42
			Maintain water infrastructure near roads (water drinking hazard)	page 42
			Maintain records of vehicle strike hotspots	page 42
4	Carnaby's Hindlimb Paralysis Syndrome	For details of this threat, see page 44	Promote awareness of this disease and what the public can do if they find a debilitated black cockatoo	page 44
5	Raven attack	For details of this threat, see page 45	Manage food waste to avoid attracting ravens	page 45
			Report sick or injured black cockatoos to helplines	page 45
6	Illegal shooting	For details of this threat, see page 46	Report injured birds to helplines	page 46
7	Climate change and Urban Heat Island effects	For details of this threat, see page 47	Reduce the City of Kwinana's greenhouse gas emissions	page 49
			Support City of Kwinana residents to reduce their emissions	page 49
			Prioritise planting of black cockatoo food trees for their evapotranspiration (urban cooling) capacity	page 49
			Support City of Kwinana residents and community groups to plant black cockatoo food trees for urban cooling	page 49
			Update the City's strategies, policies and plans as required	page 49
8	Cumulative impacts	For details of this threat, see page 50	Recognise and account for cumulative impacts in the City's planning and decision-making	page 50
			Avoid cumulative net loss of foraging habitat	page 50
			Advocate to state government for local government powers to protect significant trees on private property	page 50

Thank you to everyone at the City of Kwinana for your efforts to support the survival of Western Australia's iconic black cockatoos in the urban landscape. Murdoch University's black cockatoo research team hopes that this Conservation Action Plan is of use to the City's planners, decision-makers, land managers and other relevant personnel, to help safeguard the City of Kwinana's black cockatoos and their habitat.



Appendices

Appendix 1. Selected references and resources

Following is a list of the references cited in this Plan, and a selection of other references and resources to support the conservation of black cockatoos and their habitat. This list is not exhaustive. BirdLife Australia (Carnabys@birdlife.org.au) and DBCA are appropriate authorities to contact for specific information.

Note: hyperlinks are provided for some resources, and were functional at time of this Plan's publication.

Government black cockatoo and habitat guidelines, recovery plans and other resources

1. Department of Biodiversity, Conservation and Attractions, (n.d.). Fauna Notes: Artificial hollows for black cockatoos. file:///C:/Users/s930282j/Downloads/fauna_note_artificial_hollows_for_all_black_cockatoos-1.pdf
Advice about installing artificial hollows, including where not to install hollows in built-up areas. The above link includes a map of the Perth-Peel region which includes a red line, to the west of which DBCA advises that no artificial hollows should be installed (to reduce risks to the birds from installing artificial hollows in urban areas).
2. Department of Biodiversity, Conservation and Attractions, information on black cockatoos: www.dpaw.wa.gov.au/plants-and-animals/threatened-species-and-communities/threatened-animals/black-cockatoos
3. Department of Agriculture, Water and the Environment [now DCCEEW], (2022). *Referral guideline for 3 WA threatened black cockatoo species: Carnaby's Cockatoo, Baudin's Cockatoo and the Forest Red-tailed Black-cockatoo*. www.dcceew.gov.au/sites/default/files/documents/referral-guideline-3-wa-threatened-black-cockatoo-species-2022.pdf
See also the related DCCEEW factsheet: Black cockatoos and development in south-west Western Australia. www.dcceew.gov.au/sites/default/files/documents/black-cockatoos-development-south-west-wa-factsheet.pdf
Note: The referral guideline will be updated periodically; ensure to use the latest available version.
4. Department of Climate Change, Energy, the Environment and Water information on black cockatoos: Relevant background information on the biology and ecology of the black cockatoos is provided in the department's Species Profile and Threats (SPRAT) database.
5. Environment Protection Authority of Western Australia, (2019). *EPA Technical Advice: Carnaby's Cockatoo in Environmental Impact Assessment in the Perth and Peel Region*.

6. Western Australian Department of Parks and Wildlife, (2013). *Carnaby's cockatoo Recovery Plan*: www.dcceew.gov.au/sites/default/files/documents/carnabys-cockatoo-recovery-plan.pdf
Note: This 10-year plan is outdated and is being revised at time of writing, with another Recovery Plan expected to be available soon.
7. Western Australian Department of Parks and Wildlife, (2008). *Forest black cockatoo (Baudin's Cockatoo Calyptorhynchus baudinii and forest red-tailed black cockatoo Calyptorhynchus banksii naso) Recovery Plan* www.environment.gov.au/system/files/resources/48e4fc8c-9cb7-4c85-bc9f-6b847cf4c017/files/wa-forest-black-cockatoos-recovery-plan.pdf
Note: This plan is outdated and is being revised at time of writing, with another Recovery Plan expected to be available soon.

Murdoch University and other scientific research articles

1. Glossop, B., Clarke, K., Mitchell, D., Barrett, G., & Region, S. (2011). Methods for mapping of Carnaby's cockatoo habitat. Department of Environment and Conservation. Perth, Australia.
2. Groom, C., (2015). *Roost site fidelity and resource use by Carnaby's cockatoo, Calyptorhynchus latirostris, on the Swan Coastal Plain*, Western Australia. PhD thesis, University of Western Australia, Perth.
3. Johnstone, R. E., & Kirkby, T. (1999). Food of the forest red-tailed Black Cockatoo *Calyptorhynchus banksii naso* in south-west Western Australia. *Western Australian Naturalist*, 22, 167–178.
4. Johnstone, R. E., Kirkby, T., & Sarti, K. (2017). The distribution, status, movements and diet of the forest red-tailed black cockatoo in the south-west with emphasis on the Greater Perth Region, Western Australia. *Western Australian Naturalist*, 30, 193–219.
5. Le Roux, C. (2017). *Nocturnal roost tree, roost site and landscape characteristics of Carnaby's Black-Cockatoo (Calyptorhynchus latirostris) on the Swan Coastal Plain*. Masters thesis, Edith Cowan University.

6. Le Souëf, A., Holyoake, C., Vitali, S., & Warren, K. (2015). Presentation and prognostic indicators for free-living black cockatoos (*Calyptorhynchus spp.*) admitted to an Australian zoo veterinary hospital over 10 years. *Journal of Wildlife Diseases*, 51(2), 380–388.
7. Le Souëf, A., Vitali, S., Dawson, R., Vaughan-Higgins, R., & Warren, K. (2020). Hindlimb paralysis syndrome in wild Carnaby's cockatoos (*Calyptorhynchus latirostris*): A new threat for an endangered species. *Journal of Wildlife Diseases*, 56(3), 609–619.
8. Mawson, P. R., (1995). *The Red-capped Parrot Purpureicephalus spurius (Kuhl 1820): a pest by nature or necessity?* PhD Thesis, Department of Geography, University of Western Australia, Perth.
9. Riley, K., Warren, K.S, Armstrong, N., Yeap, L., Dawson, R., Mawson, P., Saunders, D., Cooper, C., Shephard, J. Accelerometry reveals limits to use of an energy-saving anthropogenic food source by a threatened species; a case of Carnaby's cockatoos (*Zanda latirostris*) and canola. *Ecology and Evolution*. 2023;13:e10598.
10. Robinson, A. (1960). The importance of the marri as a food source to south-western Australian birds. *The Western Australian Naturalist*, 7(5), 109–115.
11. Saunders, D. A. (1977). The effect of agricultural clearing on the breeding success of the White-tailed Black Cockatoo. *Emu*, 77(4), 180–184.
12. Saunders, D. A., Mawson, P. R., Dawson, R., Johnstone, R. E., Kirkby, T., Warren, K., ... & Garnett, S. T. (2021). Carnaby's Black-Cockatoo *Zanda latirostris*. In *The Action Plan for Australian Birds 2020* (pp. 402–407). CSIRO Publishing.
13. Saunders, D. A., & Pickup, G. (2023). A review of the taxonomy and distribution of Australia's endemic Calyptorhynchinae black cockatoos. *Australian Zoologist*. August 2023.
14. Small, G., Jimenez, I., Salzl, M., & Shrestha, P. (2020). Urban heat island mitigation due to enhanced evapotranspiration in an urban garden in Saint Paul, Minnesota, USA. *WIT Transactions on Ecology and the Environment*, 243, 39–45.
15. Williams, M. R., Yates, C. J., Saunders, D. A., Dawson, R., & Barrett, G. W. (2017). Combined demographic and resource models quantify the effects of potential land-use change on the endangered Carnaby's cockatoo (*Calyptorhynchus latirostris*). *Biological Conservation*, 210, 8–15.
- Note: See Appendix A of this research article (available a separate document from the main article) for information about recurring low-yield years for banksia; with implications with respect to providing sufficient food for black cockatoos.*

Resources from BirdLife Australia, Western Australian Museum and other specialist organisations

- BirdLife Australia. Information for councils and other land-owners and managers about plant selection and planting design for black cockatoo conservation: www.birdlife.org.au/how-to/plant-for-black-cockatoos
- BirdLife Australia information on black cockatoo food plants and dieback management: www.calameo.com/read/0041078955a20c96b51c8
The above is a printable pamphlet on choosing the best food plants for black cockatoos in your garden and school, and managing the risk of 'dieback disease' in native trees.
- BirdLife Australia. *Cocky Notes* newsletter. Biannual newsletter from BirdLife about southwest Western Australia's black cockatoos, including updates on research and conservation actions and how local councils and the public can get involved (e.g. the citizen-science Great Cocky Count). To subscribe to the *Cocky Notes* mailing list, email wa@birdlife.org.au
- BirdLife Australia. Identifying south-west black cockatoos: www.calameo.com/read/004107895ee311d26eaa4
- BirdLife International. 2022. *Zanda latirostris*. The IUCN Red List of Threatened Species 2022: e.T22684733A212974328. <https://dx.doi.org/10.2305/IUCN.UK.2022-1.RLTS.T22684733A212974328.en>
- BirdLife International (2023) Species factsheet: *Zanda baudinii*. <http://datazone.birdlife.org/species/factsheet/audins-black-cockatoo-zanda-baudinii>



- Dieback Working Group Inc.: www.dwg.org.au
Website with resources and publications regarding the relative resilience of various local native plants to dieback, as well as about how best to manage dieback in bushland.
- Kaarakin Black Cockatoo Conservation Centre: www.blackcockatooecorecovery.com
- Peck A. & Stokes V. (2019). *Black-cockatoo conservation and recovery: Guidelines for councils and land managers*. BirdLife Australia, November 2019.
The above is information for councils from BirdLife Australia.
- ReWild Perth. <https://rewildperth.com.au/>

Appendix 2. Black cockatoos from the City of Kwinana: Causes of admission to Perth Zoo Veterinary Department 2018–23

Dr Anna Le Souef

Wildlife and zoo veterinarian and *Keep Carnaby's Flying – Noolarks Forever* project officer, in collaboration with Perth Zoo Veterinary Department

Extracts of this report have been used in relevant sections of this plan. The full report is provided here for reference.

Background

To support the development of the Black Cockatoo Conservation Action Plan for the City of Kwinana, Murdoch University undertook an epidemiological analysis to identify the causes of admission to the Perth Zoo Veterinary Department (PZVD) for black cockatoos from the City of Kwinana 2018–23.

Causes of admission for black cockatoos from the City of Kwinana were vehicle strike, shooting and unknown/other causes.

Threatening processes associated with admission of black cockatoos to Perth Zoo Veterinary Department – Perth-Peel region

Vehicle strike

Vehicle strike is one of the most significant threats to black cockatoos throughout their distribution range, particularly in high-traffic areas and on roads where vehicles are travelling at speed. Vehicle strike causes traumatic injuries ranging from minor wounds and bruises to catastrophic bone fractures. Common injuries are keel fractures, wing fractures, concussions, and trauma causing internal bleeding. Many cases can be treated successfully, but a large proportion (approximately 40%) of vehicle strike cases have injuries too severe for treatment and are euthanased. Black cockatoos are at risk when feeding from vegetation and spilt grain on roadsides and median strips, and drinking from potholes in roads or puddles that have formed in roadside depressions.

Measures to mitigate the risk of vehicle strike include road signage, timely repair of potholes and other depressions in road surfaces and verges, and ensuring that plants which are black cockatoo food species (e.g. banksias and hakeas) are planted with set-backs of several metres from roads, and never on median strips (Note: see section on **Vehicle strike** in the Plan for recommended set-back distances for different road speed limits). Records of vehicle strike locations are important to maintain, as they help identify potential problem areas that may require investigation and mitigation measures.

Carnaby's hindlimb paralysis syndrome

Carnaby's hindlimb paralysis syndrome (CHiPS) has been recognised in Carnaby's cockatoos since 2012⁶⁹. To date Carnaby's cockatoos are the only species known to be affected. Carnaby's cockatoos with this syndrome are admitted to PZVD predominantly in the summer and autumn months with a weakness or paralysis in their hindlimbs; thought to be a delayed-onset neurological disease caused by pesticide exposure. Some black cockatoos with CHiPS are too sick to be treated and require euthanasia, but most can be treated with supportive care and released back to the wild. CHiPS-affected Carnaby's cockatoos tend to present to PZVD having been found in urban areas where there are key roost sites that support relatively large numbers of birds, and from areas where there are more people to notice the affected birds. Research into this syndrome is ongoing, however it is thought the birds may be being exposed to pesticides in agricultural areas outside Perth during the breeding season (spring and early summer), and, given the suspected delayed onset of this condition, are noticed in a debilitated condition when they return to their urban roosts and foraging areas in the non-breeding season. Mitigation measures for CHiPS are difficult to identify at present and require further research into the source(s) of the toxin. However, awareness of the condition is important, as many of the affected black cockatoos can be successfully rehabilitated if taken into care.

69 Le Souef A, Vitali, S, Dawson R, Vaughan-Higgins R and Warren K, 2020, Hindlimb paralysis syndrome in wild Carnaby's cockatoos (*Calyptorhynchus latirostris*): A new threat for an endangered species, *Journal of Wildlife Diseases*, 56(3), 2020, pp. 609–619

Raven attack

Raven attack is a threat to black cockatoos in urban areas which has emerged in the last decade and predominantly affects forest red-tailed black cockatoos. The observed increase in raven attacks on forest red-tailed black cockatoos in urban areas coincides with the increase in movements of this species into urban areas starting in the mid-2000s, to forage on introduced Cape Lilac; in line with declines in their main food sources (marri and jarrah) due to habitat clearing, fire and detrimental climate changes. There has also been an increase in ravens, a successful urban wildlife species, in Perth over the past 30 years.

Injuries caused by ravens tend to be around the tail end of the cockatoo, with a classic presentation being loss of tail feathers. There are sometimes wounds to the skin as well, and in some cases these wounds are too severe for treatment and require euthanasia. However, most raven-attacked black cockatoos can be treated and returned to the wild. 'Secondary' raven attacks sometimes occur when black cockatoos of any species are debilitated for another reason, such as vehicle strike or Carnaby's hindlimb paralysis syndrome. Secondary attacks further reduce these black cockatoos' chances of survival. Ravens are more numerous in areas populated by more people, because they benefit from the changes associated with urbanisation. The most successful measures for deterring ravens lie in appropriate waste management, such as self-closing bin lids. Although raven culls have occurred (with relevant lawful authority) in the past to control raven numbers in problem Perth-Peel areas, it is not clear if this is an effective measure to safeguard black cockatoos.

Displaced juveniles

This term is used to describe juvenile black cockatoos that have become separated from their parents. Sometimes this is through the well-intentioned rescue of a juvenile that was actually being cared for by parents which were not in the immediate vicinity, and sometimes it is due to injury or death of the parents. Some juveniles may also be targeted and injured by ravens; this appears to affect mainly forest red-tailed black cockatoos.

Note: while displaced juveniles are admitted to PZVD each year from across the Perth-Peel region, there were no confirmed cases from the City of Kwinana between 2018–23.


Illegal shooting

Black cockatoos are sometimes illegally shot, particularly when entering fruit or nut orchards. Black cockatoos that are found with shooting injuries have not necessarily been shot locally, but may have travelled from other areas after being injured. As of 2023, fines for shooting black cockatoos are up to \$400,000 for individuals⁷⁰. Shooting injuries can include soft tissue trauma, as well as bone fractures and internal bleeding.

Other/unknown

This category includes cockatoos that are found without apparent injuries. It is likely many are actually vehicle strike cases that were momentarily stunned or had minor concussions. Other causes include misadventure, which has included flying into poles, fences or power lines, and entanglement in string or fishing line. These cases are not numerous enough to be placed in their own category, but further highlight the risks that these endangered cockatoos face in the urban environment.

⁷⁰ As per section 150 (1) taking threatened species of the BC Act, as of 2023



City of Kwinana statistics 2018–23

A total of 9 forest red-tailed black cockatoos and 8 Carnaby’s cockatoos were admitted from City of Kwinana to the Perth Zoo Veterinary Department between 2018–2023. Main identified causes of admission were vehicle strike (both species) and shooting (forest red-tailed black cockatoos) (**Table A2.1** and **Figure A2.1**). Secondary raven attack was also reported for a forest red-tailed black cockatoo that had suffered primary vehicle strike injuries.

Vehicle strikes accounted for 10 of the cases (5 Carnaby’s cockatoos and 5 forest red-tailed black cockatoos) (**Figure A2.2**). Most were reported in the first half of the year, which is likely due to the increased presence of migratory flocks of Carnaby’s cockatoos moving onto the Swan Coastal Plain during this time of year, to spend the non-breeding season in this area.

Locations of vehicle strike cases in the City of Kwinana for 2018–23 are shown in the **Vehicle strike** section of the Plan.

There were no CHiPS cases admitted for the City of Kwinana; however, CHiPS cases may go unnoticed or unreported. There were two shooting cases admitted. Locations where these birds were recovered are shown in the **Illegal shooting** section of the Plan; however, the birds may have been shot at a different location, including potentially outside the City.




Figure A2.1 City of Kwinana black cockatoo admissions to Perth Zoo Veterinary Department 2018–23




Figure A2.2 City of Kwinana black cockatoo admissions to Perth Zoo Veterinary Department 2018–23: Vehicle strike data

Table A2.1 City of Kwinana black cockatoo admissions to Perth Zoo Veterinary Department 2018–23

Species	Vehicle strike	Shot	Other/unknown*	Total
Carnaby’s cockatoo	5	0	3	8
Forest red-tailed black cockatoo	5	2	2	9
				17

* included n=1 forest red-tailed black cockatoo injured by a golf ball strike



MU Murdoch
University

KeepCarnabysFlying.org.au

Carnabys@birdlife.org.au